




National Comprehensive  
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

# Uterine Neoplasms

**NCCN Evidence Blocks™**

Version 4.2019 — September 16, 2019



[NCCN.org](https://www.nccn.org)

[Continue](#)



**\*Nadeem R. Abu-Rustum, MD Ω/Chair**  
Memorial Sloan Kettering Cancer Center

**\*Catheryn M. Yashar, MD §/Vice Chair**  
UC San Diego Moores Cancer Center

**Sarah Bean, MD ≠**  
Duke Cancer Institute

**Kristin Bradley, MD §**  
University of Wisconsin  
Carbone Cancer Center

**Susana M. Campos, MD, MPH, MS †**  
Dana-Farber/Brigham and Women's  
Cancer Center

**Hye Sook Chon, MD Ω**  
Moffitt Cancer Center

**Christina Chu, MD Ω**  
Fox Chase Cancer Center

**David Cohn, MD Ω**  
The Ohio State University Comprehensive  
Cancer Center - James Cancer Hospital  
and Solove Research Institute

**Marta Ann Crispens, MD Ω**  
Vanderbilt-Ingram Cancer Center

**Shari Damast, MD §**  
Yale Cancer Center/  
Smilow Cancer Hospital

**Oliver Dorigo, MD, PhD Ω**  
Stanford Cancer Institute

**Christine M. Fisher, MD, MPH §**  
University of Colorado Cancer Center

**Peter Frederick, MD Ω**  
Roswell Park Cancer Institute

**David K. Gaffney, MD, PhD §**  
Huntsman Cancer Institute  
at the University of Utah

**Suzanne George, MD Ω**  
Dana-Farber/Brigham and Women's  
Cancer Center

**Robert Giuntoli II, MD Ω**  
Abramson Cancer Center at  
the University of Pennsylvania

**Ernest Han, MD, PhD Ω**  
City of Hope  
National Medical Center

**Warner K. Huh, MD Ω**  
O'Neal Comprehensive  
Cancer Center at UAB

**John R. Lurain, III, MD Ω**  
Robert H. Lurie Comprehensive Cancer  
Center of Northwestern University

**Andrea Mariani, MD Ω**  
Mayo Clinic Cancer Center

**David Mutch, MD Ω**  
Siteman Cancer Center at Barnes-  
Jewish Hospital and Washington  
University School of Medicine

**Christa Nagel, MD Ω**  
Case Comprehensive Cancer Center/  
University Hospitals Seidman Cancer  
Center and Cleveland Clinic Taussig  
Cancer Institute

**Larissa Nekhlyudov, MD, MPH †**  
Dana-Farber/Brigham and Women's  
Cancer Center

**Amanda Nickles Fader, MD Ω**  
The Sidney Kimmel Comprehensive  
Cancer Center at Johns Hopkins

**Steven W. Remmenga, MD Ω**  
Fred & Pamela Buffett Cancer Center

**R. Kevin Reynolds, MD Ω**  
University of Michigan  
Rogel Cancer Center

**Rachel Sisodia Ω**  
Massachusetts General Hospital  
Cancer Center

**Todd Tillmanns, MD Ω**  
St. Jude Children's Research Hospital/  
The University of Tennessee Health Science Center

**Stefanie Ueda, MD Ω**  
UCSF Helen Diller Family  
Comprehensive Cancer Center

**Renata Urban, MD Ω**  
Fred Hutchinson Cancer Research Center/  
Seattle Cancer Care Alliance

**Emily Wyse ¥**  
Patient Advocate

**NCCN**  
Nicole McMillian, MS

**Continue**

Ω Gynecologic oncology  
† Medical oncology  
§ Radiotherapy/Radiation oncology  
≠ Pathology  
¥ Patient advocacy  
† Internal medicine  
\*Discussion Section Writing Committee



[NCCN Uterine Neoplasms Panel Members](#)  
[NCCN Evidence Blocks Definitions \(EB-1\)](#)

**Uterine Neoplasms**  
[Uterine Neoplasms \(UN-1\)](#)

**Endometrial Carcinoma**  
[Disease Limited to the Uterus \(ENDO-1\)](#)  
[Suspected or Gross Cervical Involvement \(ENDO-2\)](#)  
[Suspected Extrauterine Disease \(ENDO-3\)](#)  
[Incompletely Surgically Staged \(ENDO-7\)](#)  
[Criteria for Considering Fertility-Sparing Options \(ENDO-8\)](#)  
[Surveillance \(ENDO-9\)](#)  
[Locoregional Recurrence \(ENDO-10\)](#)  
[High-Risk Carcinoma Histologies \(ENDO-11\)](#)

[Principles of Pathology \(ENDO-A\)](#)  
[Principles of Imaging \(ENDO-B\)](#)  
[Principles of Evaluation and Surgical Staging \(ENDO-C\)](#)  
[Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease \(ENDO-D\)](#)

**Uterine Sarcoma**  
[Diagnosed After Total Hysterectomy or Supracervical Hysterectomy ± Bilateral Salpingo-Oophorectomy \(UTSARC-1\)](#)  
[Diagnosed by Biopsy or Myomectomy \(UTSARC-1\)](#)  
[Low-Grade Endometrial Stromal Sarcoma \(ESS\) \(UTSARC-2\)](#)  
[High-Grade ESS, Undifferentiated Uterine Sarcoma \(UUS\), and Uterine Leiomyosarcoma \(UTSARC-3\)](#)  
[Surveillance \(UTSARC-4\)](#)  
[Recurrence \(UTSARC-5\)](#)

[Principles of Pathology \(UTSARC-A\)](#)  
[Principles of Imaging \(UTSARC-B\)](#)  
[Systemic Therapy for Uterine Sarcoma \(UTSARC-C\)](#)  
[Uterine Sarcoma Classification \(UTSARC-D\)](#)  
[Principles of Radiation Therapy \(UN-A\)](#)

[Staging \(ST-1\)](#)

**Clinical Trials:** NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN Member Institutions, [click here: nccn.org/clinical\\_trials/clinicians.aspx](#).

**NCCN Categories of Evidence and Consensus:** All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

**NCCN Categories of Preference:** All recommendations are considered appropriate.

See [NCCN Categories of Preference](#)

**NCCN Guidelines for Patients®**  
available at [www.nccn.org/patients](http://www.nccn.org/patients)

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Evidence Blocks™ and NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Evidence Blocks™, NCCN Guidelines, and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2019.

**NCCN EVIDENCE BLOCKS CATEGORIES AND DEFINITIONS**

5					
4					
3					
2					
1					

**E = Efficacy of Regimen/Agent**  
**S = Safety of Regimen/Agent**  
**Q = Quality of Evidence**  
**C = Consistency of Evidence**  
**A = Affordability of Regimen/Agent**

**Example Evidence Block**

5					
4	■	■		■	
3	■	■	■	■	■
2	■	■	■	■	■
1	■	■	■	■	■

**E = 4**  
**S = 4**  
**Q = 3**  
**C = 4**  
**A = 3**

**Efficacy of Regimen/Agent**

<b>5</b>	<b>Highly effective:</b> Cure likely and often provides long-term survival advantage
<b>4</b>	<b>Very effective:</b> Cure unlikely but sometimes provides long-term survival advantage
<b>3</b>	<b>Moderately effective:</b> Modest impact on survival, but often provides control of disease
<b>2</b>	<b>Minimally effective:</b> No, or unknown impact on survival, but sometimes provides control of disease
<b>1</b>	<b>Palliative:</b> Provides symptomatic benefit only

**Safety of Regimen/Agent**

<b>5</b>	<b>Usually no meaningful toxicity:</b> Uncommon or minimal toxicities; no interference with activities of daily living (ADLs)
<b>4</b>	<b>Occasionally toxic:</b> Rare significant toxicities or low-grade toxicities only; little interference with ADLs
<b>3</b>	<b>Mildly toxic:</b> Mild toxicity that interferes with ADLs
<b>2</b>	<b>Moderately toxic:</b> Significant toxicities often occur but life threatening/fatal toxicity is uncommon; interference with ADLs is frequent
<b>1</b>	<b>Highly toxic:</b> Significant toxicities or life threatening/fatal toxicity occurs often; interference with ADLs is usual and severe

**Note: For significant chronic or long-term toxicities, score decreased by 1**

**Quality of Evidence**

<b>5</b>	<b>High quality:</b> Multiple well-designed randomized trials and/or meta-analyses
<b>4</b>	<b>Good quality:</b> One or more well-designed randomized trials
<b>3</b>	<b>Average quality:</b> Low quality randomized trial(s) or well-designed non-randomized trial(s)
<b>2</b>	<b>Low quality:</b> Case reports or extensive clinical experience
<b>1</b>	<b>Poor quality:</b> Little or no evidence

**Consistency of Evidence**

<b>5</b>	<b>Highly consistent:</b> Multiple trials with similar outcomes
<b>4</b>	<b>Mainly consistent:</b> Multiple trials with some variability in outcome
<b>3</b>	<b>May be consistent:</b> Few trials or only trials with few patients, whether randomized or not, with some variability in outcome
<b>2</b>	<b>Inconsistent:</b> Meaningful differences in direction of outcome between quality trials
<b>1</b>	<b>Anecdotal evidence only:</b> Evidence in humans based upon anecdotal experience

**Affordability of Regimen/Agent (includes drug cost, supportive care, infusions, toxicity monitoring, management of toxicity)**

<b>5</b>	<b>Very inexpensive</b>
<b>4</b>	<b>Inexpensive</b>
<b>3</b>	<b>Moderately expensive</b>
<b>2</b>	<b>Expensive</b>
<b>1</b>	<b>Very expensive</b>




All staging in guideline is based on updated 2010 FIGO staging. (See ST-1, ST-2, and ST-3)

**INITIAL EVALUATION<sup>a</sup>**

- H&P
- CBC (including platelets)
- Expert pathology review with additional endometrial biopsy as clinically indicated<sup>b,c</sup>
- Imaging<sup>d</sup>
- Consider genetic evaluation (See ENDO-A)
- Consider liver function test (LFT)/renal function tests/chemistry profile
- For elderly patients with uterine cancer, see the [NCCN Guidelines for Older Adult Oncology](#)

**INITIAL CLINICAL FINDINGS<sup>c</sup>**



<sup>a</sup>Initial preoperative evaluation for known or suspected malignancy.

<sup>b</sup>Preoperative imaging and biopsy may help to identify uterine sarcomas, although biopsy sensitivity is less than for endometrial cancer. If there is suspicion of malignant mesenchymal sarcoma, fragmentation/morcellation should be avoided.

<sup>c</sup>See [Principles of Pathology for Endometrial Carcinoma \(ENDO-A\)](#) and [Principles of Pathology for Uterine Sarcoma \(UTSARC-A\)](#).

<sup>d</sup>See [Principles of Imaging for Endometrial Carcinoma \(ENDO-B\)](#) and [Principles of Imaging for Uterine Sarcoma \(UTSARC-B\)](#).

<sup>e</sup>Should be treated as a high-grade endometrial cancer.


<sup>f</sup>Also known as malignant mixed mesodermal tumor or malignant mixed Müllerian tumor and including those with either homologous or heterologous stromal elements.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**  
All recommendations are category 2A unless otherwise indicated.  
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



**INITIAL CLINICAL FINDINGS**

**PRIMARY TREATMENT**



<sup>a</sup>See (UN-1) for clarification of uterine neoplasms.

<sup>b</sup>See [Principles of Pathology \(ENDO-A\)](#).

<sup>c</sup>Minimally invasive surgery (MIS) is the preferred approach when technically feasible. See [Principles of Evaluation and Surgical Staging \(ENDO-C\)](#).

<sup>d</sup>The degree of surgical staging to assess disease status depends on preoperative and intraoperative findings. Multidisciplinary expertise is recommended. See [Principles of Evaluation and Surgical Staging \(ENDO-C\)](#).

<sup>e</sup>Ovarian preservation may be safe in select premenopausal women with early-stage endometrioid cancer. Salpingectomy is preferred.

<sup>f</sup>See [Principles of Radiation Therapy for Uterine Neoplasms \(UN-A\)](#).

<sup>g</sup>See [Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease \(ENDO-D\)](#).

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).**

**All recommendations are category 2A unless otherwise indicated.**


**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



# NCCN Guidelines Version 4.2019

## Endometrial Carcinoma

### NCCN Evidence Blocks™



<sup>a</sup>See (UN-1) for clarification of uterine neoplasms.

<sup>b</sup>See Principles of Pathology (ENDO-A).

<sup>c</sup>Minimally invasive surgery (MIS) is the preferred approach when technically feasible. <sup>g</sup>See Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease (ENDO-D).  
See Principles of Evaluation and Surgical Staging (ENDO-C).

<sup>d</sup>The degree of surgical staging to assess disease status depends on preoperative and intraoperative findings. Multidisciplinary expertise is recommended.

See Principles of Evaluation and Surgical Staging (ENDO-C).

<sup>f</sup>See Principles of Radiation Therapy for Uterine Neoplasms (UN-A).

<sup>h</sup>See Principles of Imaging (ENDO-B).

<sup>i</sup>Clear demonstration of cervical stromal involvement.

<sup>j</sup>Based on summation of conventional external-beam fractionation and low-dose-rate brachytherapy equivalent.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.

All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 4.2019

## Endometrial Carcinoma

### NCCN Evidence Blocks™



<sup>a</sup>See (UN-1) for clarification of uterine neoplasms.

<sup>b</sup>See Principles of Pathology (ENDO-A).

<sup>c</sup>Minimally invasive surgery (MIS) is the preferred approach when technically feasible. See Principles of Evaluation and Surgical Staging (ENDO-C).

<sup>d</sup>The degree of surgical staging to assess disease status depends on preoperative and intraoperative findings. Multidisciplinary expertise is recommended.

See Principles of Evaluation and Surgical Staging (ENDO-C).

<sup>f</sup>See Principles of Radiation Therapy for Uterine Neoplasms (UN-A).

<sup>g</sup>See Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease (ENDO-D).

<sup>h</sup>See Principles of Imaging (ENDO-B).

<sup>k</sup>The surgical goal is to have no measurable residual disease.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

All staging in guideline is based on updated 2010 FIGO staging. (See ST-1)

**CLINICAL FINDINGS**

**HISTOLOGIC GRADE/ADJUVANT TREATMENT<sup>f,g,m,n</sup>**

Surgically staged: Stage I<sup>d</sup> →

FIGO Stage	Histologic Grade	Adjuvant Treatment
IA	G1, G2	Observation preferred or Vaginal brachytherapy if any risk factors <sup>o,p</sup>
	G3	Vaginal brachytherapy preferred or Consider observation if no myoinvasion and no lymphovascular space invasion <sup>o</sup>
IB <sup>l</sup>	G1, G2	Vaginal brachytherapy preferred or Consider observation if no risk factors <sup>o</sup>
	G3	RT (vaginal brachytherapy and/or EBRT) ± systemic therapy <sup>q</sup>

<sup>d</sup>The degree of surgical staging to assess disease status depends on preoperative and intraoperative findings. Multidisciplinary expertise is recommended.

[See Principles of Evaluation and Surgical Staging \(ENDO-C\).](#)

<sup>f</sup>[See Principles of Radiation Therapy for Uterine Neoplasms \(UN-A\).](#)

<sup>g</sup>[See Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease \(ENDO-D\).](#)

<sup>l</sup>Consider additional imaging if not previously done.

[Principles of Pathology \(ENDO-A\).](#)

<sup>m</sup>Adjuvant therapy determinations are made on the basis of pathologic findings and type of hysterectomy.

<sup>n</sup>Initiate RT as soon as the vaginal cuff is healed, preferably no later than 12 weeks after surgery.

<sup>o</sup>Potential adverse risk factors: age ≥60 yrs, depth of invasion, and/or lymphovascular invasion (LVSI). [See Discussion](#) for additional information on adverse risk factors.

<sup>p</sup>Consider vaginal brachytherapy if one risk factor. Vaginal brachytherapy strongly suggested if two risk factors present.

<sup>q</sup>Risk factors that would lead to EBRT ± systemic therapy are: age, LVSI, and depth of myoinvasion. Risk factors are continuous variables. Risk of recurrence is higher with older age (especially >60 yrs), extensive LVSI, and deeper myoinvasion (>50%). Also, when there are more risk factors present, the risk of recurrence is higher.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

All staging in guideline is based on updated 2010 FIGO staging. (See ST-1)

**CLINICAL FINDINGS**

**HISTOLOGIC GRADE/ADJUVANT TREATMENT<sup>f,g,m,n</sup>**

Surgically staged:<sup>d</sup>  
 Stage II<sup>l,r</sup>



FIGO Stage	Histologic Grade	Adjuvant Treatment
II	G1, G2	Vaginal brachytherapy and/or EBRT <sup>s</sup>
	G3	EBRT ± vaginal brachytherapy ± systemic therapy (category 2B for systemic therapy)

[See Surveillance \(ENDO-9\)](#)

<sup>d</sup>The degree of surgical staging to assess disease status depends on preoperative and intraoperative findings. Multidisciplinary expertise is recommended.  
[See Principles of Evaluation and Surgical Staging \(ENDO-C\).](#)

<sup>f</sup>[See Principles of Radiation Therapy for Uterine Neoplasms \(UN-A\).](#)

<sup>g</sup>[See Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease \(ENDO-D\).](#)

<sup>l</sup>Consider additional imaging if not previously done. [See Principles of Imaging \(ENDO-B\).](#)

<sup>m</sup>Adjuvant therapy determinations are made on the basis of pathologic findings and type of hysterectomy.

<sup>n</sup>Initiate EBRT as soon as the vaginal cuff is healed, preferably no later than 12 weeks after surgery.

<sup>r</sup>The adverse fundal risk factors influencing therapy decisions for stage I disease ([see ENDO-4](#)), such as depth of myometrial invasion and LVSI, may also impact the choice of adjuvant therapy for stage II disease.

<sup>s</sup>Observation or vaginal brachytherapy is also an option for these patients with stage II disease who have had a radical hysterectomy with negative surgical margins and no evidence of extrauterine disease.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**


**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

All staging in guideline is based on updated 2010 FIGO staging. (See ST-1)

**CLINICAL FINDINGS**

**ADJUVANT TREATMENT<sup>f,g,m</sup>**



[See Surveillance \(ENDO-9\)](#)

<sup>d</sup>The degree of surgical staging to assess disease status depends on preoperative and intraoperative findings. Multidisciplinary expertise is recommended.

[See Principles of Evaluation and Surgical Staging \(ENDO-C\).](#)

<sup>f</sup>[See Principles of Radiation Therapy for Uterine Neoplasms \(UN-A\).](#)

<sup>g</sup>[See Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease \(ENDO-D\).](#)

<sup>m</sup>Adjuvant therapy determinations are made on the basis of pathologic findings and type of hysterectomy.

<sup>t</sup>Additional imaging if not previously done. [See Principles of Imaging \(ENDO-B\).](#)

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**


**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

All staging in guideline is based on updated 2010 FIGO staging. (See ST-1)

**CLINICAL INTRAUTERINE FINDINGS**

**ADJUVANT TREATMENT<sup>m</sup>**



[See Surveillance \(ENDO-9\)](#)

<sup>d</sup>The degree of surgical staging to assess disease status depends on preoperative and intraoperative findings. Multidisciplinary expertise is recommended. [See Principles of Evaluation and Surgical Staging \(ENDO-C\).](#)

<sup>h</sup>[See Principles of Imaging \(ENDO-B\).](#)

<sup>m</sup>Adjuvant therapy determinations are made on the basis of pathologic findings.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**  
**All recommendations are category 2A unless otherwise indicated.**  
**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**CRITERIA FOR CONSIDERING FERTILITY-SPARING OPTIONS FOR MANAGEMENT OF ENDOMETRIAL CARCINOMA (All criteria must be met)**

- Well-differentiated (grade 1) endometrioid adenocarcinoma on dilation and curettage (D&C) confirmed by expert pathology review
- Disease limited to the endometrium on MRI (preferred) or transvaginal ultrasound<sup>h</sup>
- Absence of suspicious or metastatic disease on imaging
- No contraindications to medical therapy or pregnancy
- Patients should undergo counseling that fertility-sparing option is NOT standard of care for the treatment of endometrial carcinoma

• Consultation with a fertility expert prior to therapy  
 • Genetic counseling/testing in selected patients (See UN-1)

Continuous progestin-based therapy:  
 • Megestrol  
 • Medroxyprogesterone  
 • Levonorgestrel IUD

Endometrial evaluation every 3–6 mo (either D&C or endometrial biopsy)

Complete response by 6 mo

Encourage conception<sup>v</sup> (with continued surveillance every 6 mo)

TH/BSO with staging<sup>c,d</sup> after childbearing complete or progression of disease on endometrial sampling (see ENDO-1)

Endometrial cancer present at 6–12 months<sup>h,u</sup>

TH/BSO with staging<sup>c,d</sup> (see ENDO-1)

<sup>c</sup>Minimally invasive surgery (MIS) is the preferred approach when technically feasible. See Principles of Evaluation and Surgical Staging (ENDO-C).

<sup>d</sup>The degree of surgical staging to assess disease status depends on intraoperative findings. Multidisciplinary expertise is recommended. See Principles of Evaluation and Surgical Staging (ENDO-C).

<sup>h</sup>See Principles of Imaging (ENDO-B).

<sup>u</sup>Gunderson CC, Fader AN, Carson KA, Bristow RE. Oncologic and reproductive outcomes with progestin therapy in women with endometrial hyperplasia and grade 1 adenocarcinoma: a systematic review. 2012 Gynecol Oncol;125:477-482 and Hubbs JL, Saig RM, Abaid LN, et al. Systemic and local hormone therapy for endometrial hyperplasia and early adenocarcinoma. Obstet Gynecol 2013;121:1172-1180.

<sup>v</sup>Endometrial sampling every 6 months and progestin-based therapy are recommended if patient is not in the active process of trying to conceive.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**  
 All recommendations are category 2A unless otherwise indicated.  
 Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**SURVEILLANCE**

- Physical exam every 3–6 mo for 2–3 y, then 6 mo or annually
- CA-125 if initially elevated
- Imaging as clinically indicated<sup>h</sup>
- Patient education regarding symptoms of potential recurrence, lifestyle, obesity, exercise, smoking cessation, sexual health (including vaginal dilator use and lubricants/moisturizers), nutrition counseling, potential long-term and late effects of treatment  
 (See [NCCN Guidelines for Survivorship](#) and [NCCN Guidelines for Smoking Cessation](#))

**CLINICAL PRESENTATION**

Locoregional recurrence  
 • Negative for distant metastases on radiologic imaging<sup>h</sup>

Isolated metastases

• Consider resection and/or EBRT<sup>f</sup> or Ablative therapy  
 • Consider systemic therapy<sup>g</sup> (category 2B)

Low grade or Asymptomatic or ER/PR positive

Symptomatic or Grade 2, 3 or Large volume

**THERAPY FOR RELAPSE**

[See Therapy For Relapse \(ENDO-10\)](#)

Not amenable to local treatment or Further recurrence

Treat as disseminated metastases (See below)

Hormone therapy<sup>g</sup> → If progression, systemic therapy<sup>g</sup>

Systemic therapy<sup>g</sup> ± palliative EBRT<sup>f</sup>

If progression, Best supportive care (See [NCCN Guidelines for Palliative Care](#))

<sup>h</sup>See [Principles of Imaging \(ENDO-B\)](#).

<sup>f</sup>See [Principles of Radiation Therapy for Uterine Neoplasms \(UN-A\)](#).

<sup>g</sup>See [Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease \(ENDO-D\)](#).

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 4.2019


## Endometrial Carcinoma

### NCCN Evidence Blocks™

#### CLINICAL PRESENTATION

#### THERAPY FOR RELAPSE

#### ADDITIONAL THERAPY



<sup>f</sup>See Principles of Radiation Therapy for Uterine Neoplasms (UN-A).

<sup>g</sup>See Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease (ENDO-D).

<sup>h</sup>See Principles of Imaging (ENDO-B).

<sup>w</sup>May include patients with isolated common iliac or para-aortic lymph node recurrence.


<sup>x</sup>Consider preoperative EBRT in select patients.

<sup>y</sup>Post-resection consolidation EBRT can be considered in patients who were not previously irradiated or who are deemed to have additional tolerance for radiation.

Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



All staging in guideline is based on updated 2010 FIGO staging. (See ST-1)

<sup>c</sup>Minimally invasive surgery (MIS) is the preferred approach when technically feasible. See Principles of Evaluation and Surgical Staging (ENDO-C).  
<sup>d</sup>The degree of surgical staging to assess disease status depends on intraoperative findings. Multidisciplinary expertise is recommended. See Principles of Evaluation and Surgical Staging (ENDO-C).  
<sup>f</sup>See Principles of Radiation Therapy for Uterine Neoplasms (UN-A).  
<sup>g</sup>See Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease (ENDO-D).  
<sup>h</sup>See Principles of Imaging (ENDO-B).  
<sup>z</sup>Also known as malignant mixed mesodermal tumor or malignant mixed Müllerian tumor.  
<sup>aa</sup>Observation only for select patients with no residual serous or clear cell carcinoma in the hysterectomy specimen.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.  
 All recommendations are category 2A unless otherwise indicated.  
 Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF PATHOLOGY<sup>a,1,2</sup>****Procedure:**

- TH/BSO: Total hysterectomy + bilateral salpingo-oophorectomy
- RH: Radical hysterectomy

**Pathologic assessment for carcinoma (including carcinoma, carcinosarcoma, and neuroendocrine carcinoma):**

- Uterus
  - ▶ Hysterectomy type
  - ▶ Specimen integrity (intact, opened, morcellated, other)
  - ▶ Tumor site (endometrium, lower uterine segment, polyp)
  - ▶ Tumor size
  - ▶ Histologic type
  - ▶ Histologic grade
  - ▶ Myometrial invasion (depth of invasion in mm/myometrial thickness in mm)
  - ▶ Cervical stromal involvement<sup>b</sup>
  - ▶ Lymphovascular invasion
- Other tissue/organ involvement (fallopian tubes, ovaries, vagina, parametrium, peritoneum, omentum, other)
- Peritoneal/ascitic fluid cytology<sup>c</sup>
- Lymph nodes (when resected)
  - ▶ Sentinel lymph nodes should undergo ultrastaging for detection of low-volume metastasis<sup>d</sup>
  - ▶ Level of nodal involvement (ie, pelvic, common iliac, para-aortic)
  - ▶ Number of lymph nodes with isolated tumor cells, micrometastasis, macrometastasis
- Universal testing of endometrial carcinomas for mismatch repair (MMR) proteins/microsatellite instability (MSI)
  - ▶ Testing may be performed on the hysterectomy specimen (can also be done on presurgical biopsy)
  - ▶ *MLH1* loss should be further evaluated for promoter methylation to assess epigenetic process
  - ▶ Genetic counseling and testing for all other MMR abnormalities
  - ▶ For those who are dMMR-negative or those who have not been screened, but who have strong family history of endometrial and/or colorectal cancer, genetic counseling and testing for patients is recommended. ([See Lynch syndrome/HNPCC in the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#)).
- Estrogen receptor testing is recommended in the settings of stage III, IV, and recurrent disease.
- Consider HER2 immunohistochemistry (IHC) testing (with reflex to HER2 FISH for equivocal IHC) for possible treatment of advanced stage or recurrent serous endometrial cancer.<sup>3</sup>
- Morphologic evaluation of endometrial carcinoma to determine histologic type—especially in high-grade cancers—is challenging and issues exist regarding diagnostic reproducibility.<sup>4,5</sup>
- Molecular analysis of endometrial cancer has identified four clinically significant molecular subgroups with differing clinical prognoses: *POLE* mutations, MSI-H, copy number low, and copy number high.<sup>6</sup> Ancillary studies for *POLE* mutations, MMR/MSI, and aberrant *p53* expression may be utilized to complement morphologic assessment of histologic tumor type.<sup>7</sup>

<sup>a</sup>[See Principles of Evaluation and Surgical Staging \(ENDO-C\)](#).

<sup>b</sup>Additional information including depth of invasion in mm/cervical wall thickness in mm may be requested by radiation oncologists to aid in the decision for EBRT.

<sup>c</sup>Although cytology by itself does not affect FIGO staging, cytology results should still be obtained because positive cytology is an adverse risk factor.

<sup>d</sup>Ultrastaging commonly entails thin serial sectioning of the gross sentinel lymph node and review of multiple H&E stained sections and cytokeratin immunohistochemistry for all blocks of sentinel lymph node. There is not a standard protocol for lymph node ultrastaging.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**References****ENDO-A**  
**1 OF 2**



**PRINCIPLES OF PATHOLOGY  
(REFERENCES)**

- <sup>1</sup>American College of Obstetricians and Gynecologists. ACOG practice bulletin, clinical management guidelines for obstetrician-gynecologists, number 65, August 2005: management of endometrial cancer. *Obstet Gynecol* 2005;106:413-425.
- <sup>2</sup>Krishnamurti U, Movahedi-Lankarani, S, Birdsong GG, et al. Protocol for the examination of specimens from patients with carcinoma and carcinosarcoma of the endometrium. College of American Pathologists 2018.
- <sup>3</sup>Fader AN, Rogue DM, Siegel E, et al. Randomized phase II trial of carboplatin-paclitaxel versus carboplatin-paclitaxel-trastuzumab in uterine serous carcinomas that overexpress human epidermal growth factor receptor 2/neu. *J Clin Oncol* 2018;36:2044-2051.
- <sup>4</sup>Hoang L, Kinloch MA, Leo JM, et al. Interobserver agreement in endometrial carcinoma histotype diagnosis varies depending on The Cancer Genome Atlas (TCGA)-based molecular subgroup. *Am J Surg Pathol* 2017;41(2):245-252.
- <sup>5</sup>Thomas S, Hussein Y, Bandyopadhyay S, et al. Interobserver variability in the diagnosis of uterine high-grade endometrioid carcinoma. *Arch Pathol Lab Med* 2016;140:836-843.
- <sup>6</sup>The Cancer Genome Atlas (TCGA) Research Network; Kandoth C, Schultz N, Cherniack AD, et al. Integrated genomic characterization of endometrial carcinoma. *Nature* 2013;497:67-73.
- <sup>7</sup>Murali R, Delair DF, Bean SM, et al. Evolving roles of histologic evaluation and molecular/genomic profiling in the management of endometrial cancer. *JNCCN J Natl Compr Canc Netw* 2018;16:201-209.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF IMAGING<sup>a,1-9</sup>****Initial Workup****• Non-Fertility-Sparing Treatment**

- ▶ Consider chest imaging (chest x-ray). If an abnormality is seen then chest CT without contrast may be performed.
- ▶ Consider pelvic MRI to establish the origin of the tumor (endocervical vs. endometrial) and assess local disease extent.
- ▶ For high-grade carcinoma,<sup>b</sup> consider chest/abdominal/pelvic CT to evaluate for metastatic disease.
- ▶ For patients who underwent TH with incidental finding of endometrial cancer or incompletely staged with uterine risk factors,<sup>c</sup> consider chest/abdominal/pelvic CT to evaluate for metastatic disease.
- ▶ Consider whole body PET/CT if metastasis is suspected in select patients.
- ▶ Other initial imaging should be based on symptomatology and clinical concern for metastatic disease.<sup>d</sup>

**• Fertility-Sparing Treatment**

- ▶ Pelvic MRI (preferred) to exclude myoinvasion and assess local disease extent; pelvic transvaginal ultrasound if MRI contraindicated.
- ▶ Consider chest imaging (chest x-ray). If an abnormality is seen then chest CT without contrast may be performed.
- ▶ Consider whole body PET/CT if metastasis is suspected in select patients.
- ▶ Other imaging should be based on symptomatology and clinical concern for metastatic disease.<sup>e</sup>

**Follow-up/Surveillance****• Non-Fertility-Sparing Treatment**

- ▶ Imaging should be based on symptomatology and clinical concern for metastatic disease.<sup>e</sup>
- ▶ For patients with treated FIGO stage III-IV disease optional chest/abdominal/pelvic CT every 6 months for the first 3 years and then every 6–12 months for the next 2 years.

**• Fertility-Sparing Treatment**

- ▶ Repeat pelvic MRI (preferred) for patients with persistent endometrial carcinoma after 6 months of failed medical therapy, especially if considering further fertility-sparing approaches.
- ▶ Other imaging should be based on symptomatology and clinical concern for metastatic disease.<sup>e</sup>

**Suspected Recurrence or Metastasis**

- Abdominal/pelvic and/or chest CT based upon symptoms or physical exam findings<sup>e</sup>
- Consider whole body PET/CT in select patients who may be candidates for surgery/locoregional therapy
- Consider pelvic MRI for patients who still retain their uterus

<sup>a</sup>MRI and CT are performed with contrast throughout the guidelines unless contraindicated. Contrast is not required for screening chest CT.

<sup>b</sup>High-grade endometrial carcinoma includes: poorly differentiated endometrioid, serous, clear cell, undifferentiated carcinoma, and carcinosarcoma.

<sup>c</sup>Uterine risk factors identified post TH include: high-grade carcinomas (above criteria), myoinvasion >50%, cervical stromal involvement, LVI, and tumor >2 cm.

<sup>d</sup>Indications may include abnormal physical exam findings; bulky uterine tumor; vaginal or extrauterine involvement; delay in presentation or treatment; and abdominal or pulmonary symptoms.

<sup>e</sup>Indications may include abnormal physical exam findings such as vaginal tumor; palpable mass or adenopathy; and new pelvic, abdominal, or pulmonary symptoms.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**References**



## PRINCIPLES OF IMAGING (References)

- <sup>1</sup>Salani R, Khanna N, Frimer M, et al. An update on post-treatment surveillance and diagnosis of recurrence in women with gynecologic malignancies: Society of Gynecologic Oncology (SGO) recommendations. *Gynecol Oncol* 2017;146:3-10.
- <sup>2</sup>Haldorsen IS, Salvesen HB. What is the best preoperative imaging for endometrial cancer? *Curr Oncol Rep* 2016 Apr;18(4):25.
- <sup>3</sup>Elit L, Reade CJ. Recommendations for follow-up care for gynecologic cancer survivors. *Obstet Gynecol* 2015 Dec;126 (6):1207-1214.
- <sup>4</sup>Vargas HA, Akin O, Zheng J, et al. The value of MR imaging when the site of uterine cancer origin is uncertain. *Radiology* 2011 Mar;258(3):785-792.
- <sup>5</sup>Sohaib SA, Houghton SL, Meroni R, et al. Recurrent endometrial cancer: patterns of recurrent disease and assessment of prognosis. *Clin Radiol* 2007 Jan;62(1):28-34; discussion 35-36.
- <sup>6</sup>Hensley ML, Barrette BA, Baumann K, et al. Gynecologic Cancer InterGroup (GCIG) consensus review: uterine and ovarian leiomyosarcomas. *Int J Gynecol Cancer* 2014 Nov;24(9 Suppl 3):S61-66.
- <sup>7</sup>Lakhman Y, Katz SS, Goldman DA, et al. Diagnostic performance of computed tomography for preoperative staging of patients with non-endometrioid carcinomas of the uterine corpus. *Ann Surg Oncol* 2016 Apr;23(4):1271-1278.
- <sup>8</sup>Colombo N, Creutzberg C, Amant F, et al. ESMO-ESGO-ESTRO Endometrial Consensus Conference Working Group. ESMO-ESGO-ESTRO Consensus Conference on Endometrial Cancer: diagnosis, treatment and follow-up. *Ann Oncol* 2016 Jan;27(1):16-41.
- <sup>9</sup>Sala E, Rockall AG, Freeman SJ, et al. The added role of MR imaging in treatment stratification of patients with gynecologic malignancies: what the radiologist needs to know. *Radiology* 2013 Mar;266(3):717-740.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF EVALUATION AND SURGICAL STAGING****Principles of Surgical Staging for Endometrial Cancer**<sup>1-15</sup>

- TH/BSO, and lymph node assessment is the primary treatment of apparent uterine-confined endometrial carcinoma, unless patients desire (and are candidates for) fertility-sparing options ([See ENDO-8](#)).<sup>1-3</sup> Select patients with metastatic endometrial carcinoma are also candidates for hysterectomy. ([See Principles of Pathology \[ENDO-A\]](#))
- Endometrial carcinoma should be removed en bloc to optimize outcomes; intraperitoneal morcellation or tumor fragmentation should be avoided.
- TH/BSO and lymph node assessment may be performed by any surgical route (eg, laparoscopic, robotic, vaginal, abdominal), although the standard in those with apparent uterine-confined disease is to perform the procedure via a minimally invasive approach. Randomized trials, a Cochrane Database Systematic Review, and population-based surgical studies support that minimally invasive techniques are preferred in this setting due to a lower rate of surgical site infection, transfusion, venous thromboembolism, decreased hospital stay, and lower cost of care, without compromise in oncologic outcome.<sup>4-9</sup>
- The lymph node assessment includes evaluation of the nodal basins that drain the uterus, and often comprises a pelvic nodal dissection with or without para-aortic nodal dissection. This continues to be an important aspect of surgical staging in women with uterine-confined endometrial carcinoma, as the procedure provides important prognostic information that may alter treatment decisions.
- Pelvic lymph nodes from the external iliac, internal iliac, obturator, and common iliac nodes are frequently removed for staging purposes.
- Para-aortic nodal evaluation from the inframesenteric and infrarenal regions may also be utilized for staging in women with high-risk tumors such as deeply invasive lesions, high-grade histology, and tumors of serous carcinoma, clear cell carcinoma, or carcinosarcoma.
- Sentinel lymph node (SLN) mapping may be considered. ([See pages 2–6 of ENDO-C](#))<sup>15</sup>
- Excision of suspicious or enlarged lymph nodes in the pelvic or aortic regions is important to exclude nodal metastasis.
- Some patients may not be candidates for lymph node dissection.
- Visual evaluation of the peritoneal, diaphragmatic, and serosal surfaces with biopsy of any suspicious lesions is important to exclude extrauterine disease.
- While peritoneal cytology does not impact staging, FIGO and AJCC nonetheless recommend that surgeons continue to obtain this during the TH/BSO.
- Omental biopsy is commonly performed in those with serous carcinoma, clear cell carcinoma, or carcinosarcoma histologies.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)**ENDO-C**  
**1 OF 6**

**PRINCIPLES OF EVALUATION AND SURGICAL STAGING WHEN SLN MAPPING IS USED****Principles of Sentinel Lymph Node(s) Mapping for Endometrial Cancer Staging**<sup>10-26</sup>

- Prospective and retrospective studies demonstrate that compared to systemic lymphadenectomy, SLN mapping with ultrastaging may increase the detection of lymph node metastasis with low false-negative rates in women with apparent uterine-confined disease.<sup>10-23,26</sup> If SLN mapping is considered, the expertise of the surgeon and attention to technical detail is critical. Recent evidence indicates that sentinel node mapping may also be used in high-risk histologies (serous carcinoma, clear cell carcinoma, carcinosarcoma).<sup>24,25</sup>
- SLN mapping can be considered for the surgical staging of apparent uterine-confined malignancy when there is no metastasis demonstrated by imaging studies or no obvious extrauterine disease at exploration.
- A cervical injection with dye has emerged as a useful and validated technique for identification of lymph nodes that are at high risk for metastases (ie, SLN in patients with early-stage endometrial cancer<sup>10-12</sup>).
- Superficial (1–3 mm) and optional deep (1–2 cm) cervical injection leads to dye delivery to the main layers of lymphatic channel origins in the cervix and corpus, namely the superficial subserosal, intermediate stromal, and deep submucosal lymphatic sites of origin (Figure 1 on [ENDO-C 3 of 6](#)).<sup>26,27</sup>
- Injection into the uterine cervix provides excellent dye penetration to the region of the uterine vessels and main uterine lymphatic trunks that condense in the parametria and appear in the broad ligament leading to pelvic and occasionally paraaortic sentinel nodes.
- The uterine body lymphatic trunks commonly cross over the obliterated umbilical artery with the most common location of pelvic SLN being medial to the external iliac, ventral to the hypogastric, or in the superior part of the obturator region (Figure 2 on [ENDO-C 3 of 6](#)).
- A less common location is usually seen when the lymphatic trunks do not cross over the obliterated umbilical and move cephalad following the mesoureter; in these cases, the SLN is usually seen in the common iliac presacral region (Figure 3 on [ENDO-C 3 of 6](#)).
- The radiolabeled colloid most commonly injected into the cervix is technetium-99m (99mTc); colored dyes are available in a variety of forms (Isosulfan Blue 1% and Methylene Blue 1%, Patent Blue 2.5% sodium).
- Indocyanine green (ICG) recently emerged as a useful imaging dye that requires near-infrared camera for localization, provides a very high SLN detection rate, and is commonly used in many practices at the present time.<sup>20,26</sup>
- Low-volume nodal metastasis to SLN detected only by enhanced pathologic ultrastaging is another potential value to staging with SLN.<sup>10,21-23</sup>
- The key point to a successful SLN mapping is the adherence to the SLN algorithm, which requires the performance of a side-specific nodal dissection in cases of failed mapping and removal of any suspicious or grossly enlarged nodes regardless of mapping (Figure 4 on [ENDO-C 4 of 6](#)).<sup>10-12,23,25</sup>
- For cases of failed SLN mapping, intraoperative assessment may be used to guide treatment.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)**ENDO-C**  
**2 OF 6**



## PRINCIPLES OF EVALUATION AND SURGICAL STAGING WHEN SLN MAPPING IS USED

### Principles of Sentinel Lymph Node(s) Mapping for Endometrial Cancer Staging (continued)<sup>10-26</sup>

- Sentinel lymph nodes are processed using ultrastaging, which typically includes two components:
  1. Serial sectioning with review of multiple H&E stained slides, and
  2. Cytokeratin IHC staining.
- ▶ Protocols of serial sectioning and ultrastaging vary among gynecologic pathologists.<sup>28</sup> Comparison of two different ultrastaging protocols in endometrial cancer SLN did not reveal significant advantages when serial H&E sectioning and IHC staining were used.<sup>29</sup>
- Recent data highlight the potential importance of ultrastaging for detection of low-volume metastasis. In general, SLN mapping allows for increased intraoperative surgical precision to identify nodes more likely to harbor metastasis coupled with enhanced pathology protocols, which has been shown to increase the detection of nodal metastasis, which may alter stage and adjuvant therapy recommendations.
- Lymph nodes with isolated tumor cells should be clearly reported. In endometrial cancer, when isolated tumor cells are detected in the absence of macrometastasis and micrometastasis, the lymph node stage is designated pN0(i+).<sup>30</sup>


**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF EVALUATION AND SURGICAL STAGING WHEN SLN MAPPING IS USED**


**Figure 1: Common cervical injection sites for mapping uterine cancer†**



**Figure 2: Most common location of SLNs (blue, arrow) following a cervical injection†**



**Figure 3: Less common location of SLNs (green, arrow) usually seen when lymphatic trunks are not crossing over the umbilical ligament but following the mesoureter cephalad to common iliac and presacral region†**



†Figures 1, 2, and 3 are reproduced with permission from Memorial Sloan Kettering Cancer Center. © 2013, Memorial Sloan Kettering Cancer Center.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).**

**All recommendations are category 2A unless otherwise indicated.**


**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

[Continued](#)

**ENDO-C**  
**4 OF 6**

**PRINCIPLES OF EVALUATION AND SURGICAL STAGING WHEN SLN MAPPING IS USED**

**Figure 4: The SLN algorithm for surgical staging of endometrial cancer\***



\*Reproduced with permission from Barlin JN, Khoury-Collado F, Kim CH, et al. The importance of applying a sentinel lymph node mapping algorithm in endometrial cancer staging: Beyond removal of blue nodes. *Gynecol Oncol* 2012;125:531-535.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**Continued**

**ENDO-C**  
**5 OF 6**

## PRINCIPLES OF EVALUATION AND SURGICAL STAGING REFERENCES

- <sup>1</sup>American College of Obstetricians and Gynecologists. ACOG practice bulletin, clinical management guidelines for obstetrician-gynecologists, number 65, August 2005: management of endometrial cancer. *Obstet Gynecol* 2005;106:413-425.
- <sup>2</sup>Bakkum-Gamez JN, Gonzalez-Bosquet J, Laack NN, et al. Current issues in the management of endometrial cancer. *Mayo Clin Proc* 2008 Jan;83:97-112.
- <sup>3</sup>Edge SB, Byrd DR, Compton CC. *AJCC Cancer Staging Manual*, 7th edition. New York: Springer; 2010.
- <sup>4</sup>Walker JL, Piedmonte MR, Spirtos NM, et al. Laparoscopy compared with laparotomy for comprehensive surgical staging of uterine cancer: Gynecologic Oncology Group Study LAP2. *J Clin Oncol* 2009 Nov 10;27(32):5331-6.
- <sup>5</sup>Kornblith AB, Huang HQ, Walker JL, et al. Quality of life of patients with endometrial cancer undergoing laparoscopic international federation of gynecology and obstetrics staging compared with laparotomy: a Gynecologic Oncology Group study. *J Clin Oncol* 2009;27:5337-5342.
- <sup>6</sup>Galaal K, Bryant A, Fisher AD, et al. Laparoscopy versus laparotomy for the management of early stage endometrial cancer. *The Cochrane Database of Systematic Reviews* 2012, Issue 9.
- <sup>7</sup>Scalici J, Laughlin BB, Finan MA, et al. The trend towards minimally invasive surgery (MIS) for endometrial cancer: an ACS NSQIP evaluation of surgical outcomes. *Gynecol Oncol* 2015;136:512-5.
- <sup>8</sup>Fader AN, Weise RM, Sinno AK, et al. Utilization of minimally invasive surgery in endometrial cancer care: a quality and cost disparity. *Obstet Gynecol* 2016 Jan;127(1):91-100.
- <sup>9</sup>Mannschreck D, Weise RM, Dowdy SC, et al. Disparities in surgical care among women with endometrial cancer. *Obstet Gynecol* 2016 Sept; 128:526-534.
- <sup>10</sup>Abu-Rustum NR, Khoury-Collado F, Pandit-Taskar N, et al. Sentinel lymph node mapping for grade 1 endometrial cancer: is it the answer to the surgical staging dilemma? *Gynecol Oncol* 2009;113:163-169.
- <sup>11</sup>Khoury-Collado F, Glaser GE, Zivanovic O, et al. Improving sentinel lymph node detection rates in endometrial cancer: how many cases are needed? *Gynecol Oncol* 2009;115:453-455.
- <sup>12</sup>Khoury-Collado F, Murray MP, Hensley ML, et al. Sentinel lymph node mapping for endometrial cancer improves the detection of metastatic disease to regional lymph nodes. *Gynecol Oncol* 2011;122:251-254.
- <sup>13</sup>Frimer M, Khoury-Collado F, Murray MP, et al. Micrometastasis of endometrial cancer to sentinel lymph nodes: is it an artifact of uterine manipulation? *Gynecol Oncol* 2010;119:496-499.
- <sup>14</sup>Leitao MM Jr, Khoury-Collado F, Gardner G, et al. Impact of incorporating an algorithm that utilizes sentinel lymph node mapping during minimally invasive procedures on the detection of stage IIIC endometrial cancer. *Gynecol Oncol* 2013;129:38-41.
- <sup>15</sup>Holloway RW, Abu-Rustum NR, Backes FJ, et al. Sentinel lymph node mapping and staging in endometrial cancer: A society of gynecologic oncology literature review with consensus recommendations. *Gynecol Oncol* 2017;146:405-415.
- <sup>16</sup>Kim CH, Soslow RA, Park KJ, et al. Pathologic ultrastaging improves micrometastasis detection in sentinel lymph nodes during endometrial cancer staging. *Int J Gynecol Cancer* 2013;23:964-970.
- <sup>17</sup>Barlin JN, Khoury-Collado F, Kim CH, et al. The importance of applying a sentinel lymph node mapping algorithm in endometrial cancer staging: Beyond removal of blue nodes. *Gynecol Oncol* 2012;125:531-535.
- <sup>18</sup>Vidal F, Leguevaque P, Motton S, Det al. Evaluation of the sentinel lymph node algorithm with blue dye labeling for early-stage endometrial cancer in a multicentric setting. *Int J Gynecol Cancer* 2013; 23:1327-1243.
- <sup>19</sup>Abu-Rustum NR. The Increasing credibility of sentinel lymph node mapping in endometrial cancer. *Ann Surg Oncol* 2013;20:353-354.
- <sup>20</sup>Sinno AK, Fader AN, Roche KL, et al. A comparison of colorimetric versus fluorometric sentinel lymph node mapping during robotic surgery for endometrial cancer. *Gynecol Oncol* 2014 Aug;134:281-286.
- <sup>21</sup>Holloway RW, Gupta S, Stavitski NM, et al. Sentinel lymph node mapping with staging lymphadenectomy for patients with endometrial cancer increases the detection of metastasis. *Gynecol Oncol* 2016 May;141(2):206-210.
- <sup>22</sup>Paley P, Veljovich DS, Press JZ, et al. A prospective investigation of fluorescence imaging to detect sentinel lymph nodes at robotic-assisted endometrial cancer staging. *Am J Obstet Gynecol* 2016 Jul;215(1):117.e1-7.
- <sup>23</sup>Sinno AK, Peijnenberg E, Fader AN, et al. Reducing overtreatment: a comparison of lymph node assessment strategies for endometrial cancer. *Gynecol Oncol*, In press, 2016 Aug [Epub ahead of print].
- <sup>24</sup>Schiavone MB, Zivanovic O, Zhou Q, et al. Survival of patients with uterine carcinosarcoma undergoing sentinel lymph node mapping. *Ann Surg Oncol* 2016;23:196-202.
- <sup>25</sup>Soliman PT, Westin SN, Dioun S, et al. A prospective validation study of sentinel lymph node mapping for high-risk endometrial cancer. *Gynecol Oncol* 2017;146:234-239.
- <sup>26</sup>Rossi EC, Kowalski LD, Scalici J, et al. A comparison of sentinel lymph node biopsy to lymphadenectomy for endometrial cancer staging (FIRES trial): a multicentre, prospective, cohort study. *Lancet Oncol* 2017;18:384-392.
- <sup>27</sup>Rossi EC, Kowalski LD, Scalici J, et al. A comparison of sentinel lymph node biopsy to lymphadenectomy for endometrial cancer staging (FIRES trial): a multicentre, prospective, cohort study. *Lancet Oncol* 2017;18:384-392.
- <sup>28</sup>Cormier B, Rozenholc AT, Gotlieb W, et al. Sentinel lymph node procedure in endometrial cancer: A systematic review and proposal for standardization of future research. *Gynecol Oncol* 2015 Aug;138:478-485.
- <sup>29</sup>Euscher E, Sui D, Soliman P, et al. Ultrastaging of Sentinel Lymph Nodes in Endometrial Carcinoma According to Use of 2 Different Methods. *Int J Gynecol Pathol* 2018 May;37:242-251.
- <sup>30</sup>Olawaiye AB, Mutch DG. Lymphnode staging update in the American Joint Committee on Cancer 8th Edition cancer staging manual. *Gynecol Oncol* 2018;150:7-8.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



**SYSTEMIC THERAPY FOR RECURRENT, METASTATIC, OR HIGH-RISK DISEASE**  
(Strongly encourage participation in clinical trials)

<p><b><u>Chemotherapy Regimens<sup>a,b</sup></u></b> <b><u>Preferred Regimens</u></b></p> <ul style="list-style-type: none"> <li>• Carboplatin/paclitaxel<sup>1</sup></li> <li>• Carboplatin/paclitaxel/trastuzumab (for HER2-positive uterine serous carcinoma)<sup>2,c</sup></li> </ul> <p><b><u>Other Recommended Regimens</u></b></p> <ul style="list-style-type: none"> <li>• Carboplatin/docetaxel<sup>d</sup></li> <li>• Cisplatin/doxorubicin<sup>3</sup></li> <li>• Cisplatin/doxorubicin/paclitaxel<sup>3,e</sup></li> <li>• Carboplatin/paclitaxel/bevacizumab<sup>4,f,g</sup></li> <li>• Ifosfamide/paclitaxel (category 1 for carcinosarcoma)<sup>5</sup></li> <li>• Cisplatin/ifosfamide (for carcinosarcoma)</li> <li>• Everolimus/letrozole (for endometrioid histology)</li> </ul> <p><b><u>Useful In Certain Circumstances</u></b></p> <ul style="list-style-type: none"> <li>• Pembrolizumab<sup>j</sup> (for MSI-H/dMMR tumors)</li> </ul>	<p><b><u>Adjuvant Treatment When Used for Uterine-Confined Disease</u></b> <b><u>Preferred Regimens</u></b></p> <ul style="list-style-type: none"> <li>• Carboplatin/paclitaxel</li> </ul> <p><b><u>Hormone Therapy<sup>k</sup></u></b> <b><u>Preferred Regimens</u></b></p> <ul style="list-style-type: none"> <li>• Medroxyprogesterone acetate/tamoxifen (alternating)</li> <li>• Megestrol acetate/tamoxifen (alternating)</li> <li>• Progestational agents             <ul style="list-style-type: none"> <li>▶ Medroxyprogesterone acetate</li> <li>▶ Megestrol acetate</li> <li>▶ Levonorgestrel intrauterine device (IUD) (for select fertility-sparing cases)</li> </ul> </li> <li>• Aromatase inhibitors</li> <li>• Tamoxifen</li> <li>• Fulvestrant</li> </ul> <p>See Evidence Blocks on <a href="#">ENDO-D (EB-1)</a> and <a href="#">ENDO-D (EB-2)</a></p>
--	---

<sup>a</sup>Cisplatin, carboplatin, liposomal doxorubicin, paclitaxel, and docetaxel may cause drug reactions. (See [NCCN Guidelines for Ovarian Cancer--Management of Drug Reactions \[OV-C\]](#))

<sup>b</sup>Chemotherapy regimens can be used for all carcinoma histologies. Carcinosarcomas are now considered and treated as high-grade carcinomas. However, ifosfamide-based regimens were previously used for carcinosarcomas.

<sup>c</sup>Only for advanced (stage III/IV) and recurrent uterine serous carcinomas that are HER2-positive.

<sup>d</sup>Docetaxel may be considered for patients in whom paclitaxel is contraindicated.

<sup>e</sup>The cisplatin/doxorubicin/paclitaxel regimen is not widely used because of concerns about toxicity.

<sup>f</sup>For advanced and recurrent disease only.

<sup>g</sup>An FDA-approved biosimilar is an appropriate substitute for bevacizumab.

<sup>h</sup>Albumin-bound paclitaxel is a reasonable substitute for patients with a hypersensitivity to paclitaxel if the skin testing to paclitaxel is negative. If the patient has a positive skin test to paclitaxel then the patient requires desensitization to paclitaxel. Albumin-bound paclitaxel is not a reasonable substitute for paclitaxel if the patient's skin test is positive.

<sup>i</sup>Bevacizumab may be considered for use in patients who have progressed on prior cytotoxic chemotherapy.

<sup>j</sup>For recurrent endometrial cancer, NCCN recommends MSI-H or dMMR testing if not previously done. Pembrolizumab is indicated for patients with MSI-H or dMMR tumors that have progressed following prior cytotoxic chemotherapy.

<sup>k</sup>Hormonal therapy is typically used for lower-grade endometrioid histologies, preferably in patients with small tumor volume or an indolent growth pace.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**[References](#)**



5					E = Efficacy of Regimen/Agent
4					S = Safety of Regimen/Agent
3					Q = Quality of Evidence
2					C = Consistency of Evidence
1					A = Affordability of Regimen/Agent
	E	S	Q	C	A

**EVIDENCE BLOCKS FOR SYSTEMIC THERAPY FOR RECURRENT, METASTATIC, OR HIGH-RISK DISEASE**  
(Strongly encourage participation in clinical trials)

ADJUVANT THERAPY FOR UTERINE-CONFINED ENDOMETRIAL CARCINOMA	
Preferred Regimens	
Carboplatin/paclitaxel	

CHEMOTHERAPY REGIMENS	
Preferred Regimens	
Carboplatin/paclitaxel	
Carboplatin/paclitaxel/trastuzumab (for HER2-positive uterine serous carcinoma)	
Other Recommended Regimens	
Albumin-bound paclitaxel	
Bevacizumab	
Carboplatin	
Carboplatin/docetaxel	
Carboplatin/paclitaxel/bevacizumab	
Cisplatin	
Cisplatin/doxorubicin	

CHEMOTHERAPY REGIMENS	
Other Recommended Regimens (continued)	
Cisplatin/doxorubicin/paclitaxel	
Cisplatin/ifosfamide (for carcinosarcoma)	
Docetaxel	
Doxorubicin	
Everolimus/letrozole	
Ifosfamide (for carcinosarcoma)	
Ifosfamide/paclitaxel (for carcinosarcoma)	
Lenvatinib/pembrolizumab	*
Liposomal doxorubicin	
Paclitaxel	
Temsirolimus	
Topotecan	
Useful In Certain Circumstances	
Pembrolizumab (for MSI-H/dMMR tumors)	

\*Evidence Block development in progress

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**  
All recommendations are category 2A unless otherwise indicated.  
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued on ENDO-D \(EB-2\)](#)

**ENDO-D  
EB-1**



5						E = Efficacy of Regimen/Agent
4						S = Safety of Regimen/Agent
3						Q = Quality of Evidence
2						C = Consistency of Evidence
1						A = Affordability of Regimen/Agent
	E	S	Q	C	A	

**EVIDENCE BLOCKS FOR SYSTEMIC THERAPY FOR RECURRENT, METASTATIC, OR HIGH-RISK DISEASE**  
(Strongly encourage participation in clinical trials)

<b>HORMONE THERAPY</b>	
<b>Preferred Regimens</b>	
<b>Aromatase inhibitors</b>	
<b>Fulvestrant</b>	
<b>Medroxyprogesterone acetate/tamoxifen (alternating)</b>	
<b>Megestrol/tamoxifen (alternating)</b>	
<b>Progestational agents</b>	
<b>Tamoxifen</b>	

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



**SYSTEMIC THERAPY FOR RECURRENT, METASTATIC, OR HIGH-RISK DISEASE  
(REFERENCES)**

- <sup>1</sup>Miller D, Filiaci V, Fleming G, et al. Randomized phase III noninferiority trial of first line chemotherapy for metastatic or recurrent endometrial carcinoma: a Gynecologic Oncology Group study [abstract]. *Gynecol Oncol* 2012;125:771.
- <sup>2</sup>Fader AN, Roque DM, Siegel E, et al. Randomized phase II trial of carboplatin-paclitaxel versus carboplatin-paclitaxel-trastuzumab in uterine serous carcinomas that overexpress human epidermal growth factor receptor 2/neu. *J Clin Oncol* 2018;36:2044-2051.
- <sup>3</sup>Homesley HD, Filiaci V, Gibbons SK, et al. A randomized phase III trial in advanced endometrial carcinoma of surgery and volume directed radiation followed by cisplatin and doxorubicin with or without paclitaxel: A Gynecologic Oncology Group study. *Gynecol Oncol* 2009;112:543-552.
- <sup>4</sup>Rose PG, Ali S, Moslemi-Kebria M, Simpkins F. Paclitaxel, carboplatin, and bevacizumab in advanced and recurrent endometrial carcinoma. *Int J Gynecol Cancer* 2017 Mar;27:452-458.
- <sup>5</sup>Homesley HD, Filiaci V, Markman M, et al. Phase III trial of ifosfamide with or without paclitaxel in advanced uterine carcinosarcoma: a Gynecologic Oncology Group Study. *J Clin Oncol* 2007;25:526-531.
- <sup>6</sup>Picard M, Pur L, Caiado J, et al. Risk stratification and skin testing to guide re-exposure in taxane-induced hypersensitivity reactions. *J Allergy Clin Immunol* 2016;137(4):1154-1164.
- <sup>7</sup>Aghajanian C, Sill MW, Darcy KM, et al. Phase II trial of bevacizumab in recurrent or persistent endometrial cancer: a Gynecologic Oncology Group study. *J Clin Oncol* 2011;29:2259-2265.
- <sup>8</sup>Oza AM, Elit L, Tsao MS, et al. Phase II study of temsirolimus in women with recurrent or metastatic endometrial cancer: a trial of the NCIC Clinical Trials Group. *J Clin Oncol* 2011;29:3278-3285.
- <sup>9</sup>Makker V, Rasco D, Vogelzang NJ, et al. Lenvatinib plus pembrolizumab in patients with advanced endometrial cancer: an interim analysis of a multicentre, open-label, single-arm, phase 2 trial. *Lancet Oncol*. 2019;20:711-718.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 4.2019

## Uterine Sarcoma

### NCCN Evidence Blocks™



<sup>a</sup>Preoperative imaging and biopsy may help to identify uterine sarcomas, although biopsy sensitivity is less than for endometrial cancer. If there is suspicion of malignant mesenchymal sarcoma, fragmentation/morcellation should be avoided.

<sup>b</sup>See [Principles of Pathology \(UTSARC-A\)](#).

<sup>c</sup>See [Principles of Imaging \(UTSARC-B\)](#).

<sup>d</sup>Oophorectomy individualized for reproductive-age patients. Favor BSO if ER/PR positive.

<sup>e</sup>For incidental finding of uterine sarcoma after TH/BSO or fragmented specimen: Recommend imaging and consider additional surgical resection on an individual basis.

<sup>f</sup>Uterine sarcoma should be removed *en bloc* to optimize outcomes; morcellation should be avoided.

<sup>g</sup>See [Principles of Radiation Therapy for Uterine Neoplasms \(UN-A\)](#).

<sup>h</sup>See [Systemic Therapy \(UTSARC-C\)](#).

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).**


**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



**PATHOLOGIC FINDINGS/  
HISTOLOGIC GRADE<sup>i</sup>**

**ADDITIONAL THERAPY**



<sup>g</sup>See Principles of Radiation Therapy for Uterine Neoplasms (UN-A).

<sup>h</sup>See Systemic Therapy (UTSARC-C).

<sup>i</sup>See Uterine Sarcoma Classification (UTSARC-D).


**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PATHOLOGIC FINDINGS/  
HISTOLOGIC GRADE<sup>i</sup>**

**ADDITIONAL THERAPY**



<sup>g</sup>See Principles of Radiation Therapy for Uterine Neoplasms (UN-A).

<sup>h</sup>See Systemic Therapy (UTSARC-C).

<sup>i</sup>See Uterine Sarcoma Classification (UTSARC-D).

<sup>j</sup>Observation may be an option in select, completely resected cases with no evidence of disease on postoperative imaging.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**SURVEILLANCE**

- H&P exam every 3–4 mo for 2–3 y, then every 6–12 mo
- Imaging<sup>c</sup>
- Patient education regarding symptoms of potential recurrence, lifestyle, obesity, exercise, nutrition, sexual health (including vaginal dilator use and lubricants/moisturizers), smoking cessation, nutrition counseling and potential long-term and late effects of treatment ([See NCCN Guidelines for Survivorship and NCCN Guidelines for Smoking Cessation](#))

**RECURRENCE**

**Local recurrence:**  
 • Vagina/pelvis  
 • Imaging negative for distant metastatic disease<sup>c</sup>

**Isolated metastases**

**Disseminated disease**

**THERAPY FOR RELAPSE**

[See Therapy For Relapse \(UTSARC-5\)](#)

**Resectable**

- Surgical resection or other local ablative therapy:<sup>j</sup>  
 Consider postoperative systemic therapy<sup>h</sup>  
 ▶ Consider postoperative EBRT<sup>g</sup>

**Unresectable**

Systemic therapy<sup>h</sup> and/or Local therapy (EBRT<sup>g</sup> or local ablative therapy) → If response, consider surgery

Systemic therapy<sup>h</sup> ± palliative EBRT<sup>g</sup> or Best supportive care

<sup>c</sup>See Principles of Imaging (UTSARC-B).

<sup>g</sup>See Principles of Radiation Therapy for Uterine Neoplasms (UN-A).


<sup>h</sup>See Systemic Therapy (UTSARC-C).

<sup>j</sup>Observation may be an option in select, completely resected cases with no evidence of disease on postoperative imaging.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**  
 All recommendations are category 2A unless otherwise indicated.  
 Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**RECURRENCE**

**THERAPY FOR RELAPSE**



<sup>c</sup>See Principles of Imaging (UTSARC-B).

<sup>g</sup>See Principles of Radiation Therapy for Uterine Neoplasms (UN-A).

<sup>k</sup>The use of preoperative EBRT would preclude postoperative EBRT.

<sup>l</sup>For low-grade ESS, the first choice of systemic therapy is estrogen blockade. See Systemic Therapy (UTSARC-C).

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page EB-1.**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



## PRINCIPLES OF PATHOLOGY<sup>a,1,2</sup>

### **Procedure:**

- **TH/BSO: Total hysterectomy + bilateral salpingo-oophorectomy**
- **RH: Radical hysterectomy**

### **Pathologic assessment for sarcoma (including leiomyosarcoma, adenosarcoma, endometrial stromal sarcoma, and undifferentiated sarcoma):**

- **Uterus**
  - ▶ **Hysterectomy type**
  - ▶ **Specimen integrity (intact, opened, morcellated, other)**
  - ▶ **Tumor size**
  - ▶ **Myometrial invasion (for adenosarcoma only)**
  - ▶ **Histologic type**
  - ▶ **Histologic grade (for adenosarcoma only)**
  - ▶ **Lymphovascular invasion**
- **Other tissue/organ involvement (fallopian tubes, ovaries, vagina, parametrium, peritoneum, omentum, other)**
- **Peritoneal/ascitic fluid cytology<sup>b</sup>**
- **Lymph nodes (when resected)**
  - ▶ **Level of nodal involvement (ie, pelvic, common iliac, para-aortic)**
  - ▶ **Number of lymph nodes with metastasis**

### Footnotes

<sup>a</sup>Also see [Principles of Evaluation and Surgical Staging \(ENDO-C\)](#).

<sup>b</sup>Although cytology by itself does not affect FIGO staging, cytology results should still be obtained because positive cytology is an adverse risk factor.

### References

- <sup>1</sup>American College of Obstetricians and Gynecologists. ACOG practice bulletin, clinical management guidelines for obstetrician-gynecologists, number 65, August 2005: management of endometrial cancer. *Obstet Gynecol* 2005;106:413-425.
- <sup>2</sup>Krishnamurti U, Movahedi-Lankarani S, Bell DA, et al. Protocol for the examination of specimens from patients with primary sarcoma of the uterus. *College of American Pathologists* 2018.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF IMAGING<sup>a,1-9</sup>****Initial Workup**

- Chest/abdomen/pelvic CT.
- For patients who underwent TH with incidental finding of uterine sarcoma or incompletely resected uterus/adnexa (ie, supracervical hysterectomy (SCH), myomectomy, possible tumor fragmentation, intraperitoneal morcellation) perform chest/abdominal/pelvic CT or abdominal/pelvic MRI and chest CT without contrast to evaluate for metastatic disease.
- Consider pelvic MRI to evaluate local tumor extension or residual abnormality in cases where the uterus or adnexa were not resected or incompletely resected (ie, SCH, myomectomy, possible tumor fragmentation, intraperitoneal morcellation).
- Consider whole body PET/CT to clarify ambiguous findings.
- Other imaging should be based on symptomatology and clinical concern for metastatic disease.<sup>b</sup>

**Follow-up/Surveillance**

- Chest/abdominal/pelvic CT every 3–6 months for the first 3 years and then every 6–12 months for the next 2 years. Depending on histology grade and initial stage, consider annual to bi-annual imaging thereafter up to an additional 5 years.<sup>c</sup>
- Optional abdominal/pelvic MRI and chest CT without contrast every 3–6 months for the first 3 years and then every 6–12 months for the next 2 years. Depending on histology, grade, and initial stage, consider annual to bi-annual imaging thereafter up to an additional 5 years.<sup>c</sup>
- Consider whole body PET/CT if metastasis is suspected in select patients.
- Imaging should be based on symptomatology and clinical concern for metastatic disease.<sup>d</sup>

<sup>a</sup>MRI and CT are performed with contrast throughout the guidelines unless contraindicated. Contrast is not required for screening chest CT.

<sup>b</sup>Indications may include abnormal physical exam finding; bulky uterine tumor; vaginal or extrauterine involvement; delay in presentation or treatment; and abdominal or pulmonary symptoms.

<sup>c</sup>Follow-up imaging may be as frequent as every 3 months or change based on histology grade and/or stage of tumor.

<sup>d</sup>Indications may include abnormal physical exam findings such as vaginal involvement; palpable mass or adenopathy; and new pelvic, abdominal, or pulmonary symptoms.

**Note: For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).**

**All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

[Continued](#)**UTSARC-B**  
**1 OF 2**



**PRINCIPLES OF IMAGING  
(REFERENCES)**

- <sup>1</sup>Salani R, Khanna N, Frimer M, et al. An update on post-treatment surveillance and diagnosis of recurrence in women with gynecologic malignancies: Society of Gynecologic Oncology (SGO) recommendations. *Gynecol Oncol* 2017;146:3-10.
- <sup>2</sup>Haldorsen IS, Salvesen HB. What is the best preoperative imaging for endometrial cancer? *Curr Oncol Rep* 2016 Apr;18(4):25.
- <sup>3</sup>Elit L, Reade CJ. Recommendations for follow-up care for gynecologic cancer survivors. *Obstet Gynecol* 2015 Dec;126 (6):1207-1214.
- <sup>4</sup>Vargas HA, Akin O, Zheng J, et al. The value of MR imaging when the site of uterine cancer origin is uncertain. *Radiology* 2011 Mar;258(3):785-792.
- <sup>5</sup>Sohaib SA, Houghton SL, Meroni R, et al. Recurrent endometrial cancer: patterns of recurrent disease and assessment of prognosis. *Clin Radiol* 2007 Jan;62(1):28-34; discussion 35-36.
- <sup>6</sup>Hensley ML, Barrette BA, Baumann K, et al. Gynecologic Cancer InterGroup (GCIG) consensus review: uterine and ovarian leiomyosarcomas. *Int J Gynecol Cancer* 2014 Nov;24(9 Suppl 3):S61-66.
- <sup>7</sup>Lakhman Y, Katz SS, Goldman DA, et al. Diagnostic performance of computed tomography for preoperative staging of patients with non-endometrioid carcinomas of the uterine corpus. *Ann Surg Oncol* 2016 Apr;23(4):1271-1278.
- <sup>8</sup>Colombo N, Creutzberg C, Amant F, et al. ESMO-ESGO-ESTRO Endometrial Consensus Conference Working Group. ESMO-ESGO-ESTRO Consensus Conference on Endometrial Cancer: diagnosis, treatment and follow-up. *Ann Oncol* 2016 Jan;27(1):16-41.
- <sup>9</sup>Sala E, Rockall AG, Freeman SJ, et al. The added role of MR imaging in treatment stratification of patients with gynecologic malignancies: what the radiologist needs to know. *Radiology* 2013 Mar;266(3):717-740.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



**SYSTEMIC THERAPY FOR UTERINE SARCOMA<sup>1</sup>**  
**(Clinical trials strongly recommended)**

Systemic Therapy	Hormone Therapy for Low-grade ESS or Hormone Receptor-Positive (ER/PR) uLMS <sup>4</sup>
<p><b>Preferred Regimens</b></p> <ul style="list-style-type: none"> <li>• Doxorubicin</li> <li>• Docetaxel/gemcitabine</li> </ul> <p><b>Other Recommended Regimens</b></p> <ul style="list-style-type: none"> <li>• Doxorubicin/ifosfamide</li> <li>• Doxorubicin/dacarbazine</li> <li>• Gemcitabine/dacarbazine</li> <li>• Gemcitabine/vinorelbine</li> <li>• Dacarbazine</li> <li>• Gemcitabine</li> <li>• Epirubicin</li> <li>• Ifosfamide</li> <li>• Liposomal doxorubicin</li> <li>• Pazopanib<sup>2</sup></li> <li>• Temozolomide<sup>2</sup></li> <li>• Trabectedin<sup>3</sup></li> <li>• Eribulin (category 2B)<sup>2</sup></li> </ul>	<p><b>Preferred Regimens</b></p> <ul style="list-style-type: none"> <li>• Aromatase inhibitors for low-grade ESS</li> </ul> <p><b>Other Recommended Regimens</b></p> <ul style="list-style-type: none"> <li>• Aromatase inhibitors (for ER/PR-positive uLMS)</li> <li>• Fulvestrant</li> <li>• Megestrol acetate (category 2B for ER/PR-positive uLMS)</li> <li>• Medroxyprogesterone acetate (category 2B for ER/PR-positive uLMS)</li> <li>• GnRH analogs (category 2B for low-grade ESS and ER/PR-positive uLMS)</li> </ul>

[See Evidence Blocks on UTSARC-C \(EB-1\)](#)

<sup>1</sup>See [NCCN Guidelines for Ovarian Cancer--Management of Drug Reactions \[OV-C\]](#).

<sup>2</sup>Pazopanib, temozolomide, and eribulin may be considered for use in patients with recurrent or metastatic disease who have progressed on prior cytotoxic chemotherapy.

<sup>3</sup>For uLMS that has been treated with a prior anthracycline-containing regimen.

<sup>4</sup>These hormonal therapies may be considered for patients with uLMS that is ER/PR-positive, preferably with small tumor volume or an indolent growth pace.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Back to Recurrence \(UTSARC-4\)](#)

**UTSARC-C**

5						E = Efficacy of Regimen/Agent
4						S = Safety of Regimen/Agent
3						Q = Quality of Evidence
2						C = Consistency of Evidence
1						A = Affordability of Regimen/Agent
	E	S	Q	C	A	

**EVIDENCE BLOCKS FOR SYSTEMIC THERAPY FOR UTERINE SARCOMA**  
(Clinical trials strongly recommended)

SYSTEMIC THERAPY	
Preferred Regimens	
Doxorubicin	
Docetaxel/gemcitabine	
Other Recommended Regimens	
Doxorubicin/ifosfamide	
Doxorubicin/dacarbazine	
Gemcitabine/dacarbazine	
Gemcitabine/vinorelbine	
Dacarbazine	
Gemcitabine	
Epirubicin	
Ifosfamide	
Liposomal doxorubicin	
Pazopanib	

SYSTEMIC THERAPY	
Other Recommended Regimens (continued)	
Temozolomide	
Trabectedin	
Eribulin	

HORMONE THERAPY FOR LOW-GRADE ESS OR HORMONE RECEPTOR-POSITIVE (ER/PR) uLMS	
Preferred Regimens	
Aromatase inhibitors for low-grade ESS	
Other Recommended Regimens	
Aromatase inhibitors (for ER/PR-positive uLMS)	
Fulvestrant	
Megestrol acetate	
Medroxyprogesterone acetate	
GnRH analogs	

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).  
All recommendations are category 2A unless otherwise indicated.  
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



## UTERINE SARCOMA CLASSIFICATION<sup>1</sup>

### Most Common Types of Uterine Sarcoma:

- Low-grade endometrial stromal sarcoma (ESS)<sup>2</sup>
- High-grade ESS<sup>3</sup>
- Undifferentiated uterine sarcoma (UUS)<sup>4</sup>
- Uterine leiomyosarcoma (uLMS)<sup>5</sup>

### Rare subtypes of sarcoma that can occur in the uterus:

(see the [NCCN Guidelines for Soft Tissue Sarcoma](#))

- Adenosarcomas
- PEComas
- Rhabdomyosarcoma

<sup>1</sup>Kurman RJ, Carcangiu ML, Herrington CS, Young RH. WHO Classification of Tumours of the Female Reproductive Organs, Volume 6, 2014.

<sup>2</sup>Low-grade endometrial stromal sarcomas (LGESS) are characterized by small cells with low-grade cytology and features resembling stromal cells in proliferative endometrium. Mitotic activity is usually low (<5 MF per 10 HPF).

<sup>3</sup>High-grade endometrial stromal sarcomas (HGESS) are characterized by small cells with high-grade cytology, frequent necrosis, and brisk mitotic activity (>10 MF per 10 HPF). HGESS can contain areas of conventional LGESS.

<sup>4</sup>Undifferentiated uterine sarcomas (UUSs) are characterized by cells with high-grade cytologic features lacking any resemblance to the stromal cells in proliferative endometrium or any other specific type of differentiation.

<sup>5</sup>Excludes smooth muscle tumors of uncertain malignant potential, epithelioid smooth muscle tumors, benign metastasizing leiomyomas, intravenous leiomyomatosis, and diffuse leiomyomatosis; management in individual cases may be modified based on clinicopathologic prognostic factors, such as size (< or > 5 cm), mitotic activity (< or > 10 mf/10 hpf), age (< or > 50 years), and presence or absence of vascular invasion.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



## PRINCIPLES OF RADIATION THERAPY FOR UTERINE NEOPLASMS

### General Principles-Uterine Neoplasms

- RT is directed at sites of known or suspected tumor involvement and may include EBRT and/or brachytherapy. Imaging is required to assess locoregional extent and to rule out distant metastases before administration of RT. In general, EBRT is directed to the pelvis with or without the para-aortic region. Brachytherapy can be delivered: 1) to an intact uterus, either preoperatively or definitively; or 2) more commonly, to the vagina after hysterectomy. For the purposes of these guidelines, whole abdominal radiotherapy is not considered to be tumor-directed RT.
- Evidence supports the use of combined modality radiation and chemotherapy as adjuvant treatment for patients with extrauterine disease.<sup>1</sup>

<sup>1</sup>Klopp A, Smith BD, Alektiar K, et al. The role of postoperative radiation therapy for endometrial cancer: executive summary of an american society for radiation oncology evidence-based guideline. Pract Radiat Oncol 2014;4:137-144.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF RADIATION THERAPY FOR UTERINE NEOPLASMS****General Treatment Information****• Target Volumes**

- ▶ Pelvic radiotherapy should target the gross disease (if present), the lower common iliacs, external iliacs, internal iliacs, obturators, parametria, upper vagina/para-vaginal tissue, and presacral lymph nodes (in patients with cervical involvement).
- ▶ Extended-field radiotherapy should include the pelvic volume and also target the entire common iliac chain and para-aortic lymph node region. The upper border of the extended field depends on the clinical situation but should at least be 1–2 cm above the level of the renal vessels.
- ▶ Pelvic tissues at risk, especially in the post-hysterectomy setting, can be highly variable depending on bowel and bladder filling. In this situation, the integrated target volume (ITV), which encompasses the range of organ movement and deformation, is considered the clinical target volume (CTV), and should be fully covered in the treatment volume.

**• Dosing Prescription Regimen – External Beam**

- ▶ External-beam doses for microscopic disease should be 45–50 Gy. Multiple conformal fields based on CT-treatment planning should be utilized, and consideration for IMRT for normal tissue sparing may be considered, with appropriate attention to QA and tissue interfraction mobility. Postoperatively, if there is gross residual disease and the area(s) can be sufficiently localized, a boost can be added to a total dose of 60–70 Gy, respecting normal tissue sensitivity.
- ▶ For neoadjuvant radiation, doses of 45–50 Gy are typically used. One could consider adding 1–2 high dose-rate (HDR) insertions to a total dose of 75–80 Gy low-dose-rate equivalent, to minimize risk of positive or close margins at hysterectomy.

**• Dosing Prescription Regimen – Brachytherapy**

- ▶ Initiate brachytherapy as soon as the vaginal cuff is healed, preferably 6–8 weeks after surgery but in general initiation of brachytherapy should not exceed 12 weeks. For vaginal brachytherapy, the dose should be prescribed to the vaginal surface or at a depth of 0.5 cm from the vaginal surface; the dose depends on the use of EBRT. The target for vaginal brachytherapy after hysterectomy should be no more than the upper two-thirds of the vagina; in cases of extensive LVSI or positive margins, a longer segment of the vagina may be treated.
  - ◊ For postoperative high-dose-rate (HDR) vaginal brachytherapy alone, regimens include 6 Gy x 5 fractions prescribed to the vaginal surface, or 7 Gy x 3 fractions or 5.5 Gy x 4 fractions prescribed to 5 mm below the vaginal surface. While 7 Gy x 3 fractions prescribed at a depth of 0.5 cm from the vaginal surface is a regimen used by many, the use of smaller fraction sizes may be considered to potentially further limit toxicity in selected cases.
  - ◊ When HDR brachytherapy is used as a boost to EBRT, doses of 4–6 Gy x 2 to 3 fractions prescribed to the vaginal mucosa are commonly used.
- ▶ For medically inoperable uterine cancer, risk of extrauterine spread determines the combination of external beam plus brachytherapy or brachytherapy alone. Brachytherapy doses for definitive therapy are individualized based on the clinical situation. When available, image-guided therapy should be used. Based on the best available evidence, an EQD2 D90 of at least 48 Gy should be delivered to the uterus, cervix, and upper 1–2 cm of vagina if brachytherapy alone is used, and should be increased to 65 Gy for the combination of EBRT and brachytherapy. If an MRI is used as part of planning, the target dose for the GTV would be an EQD2 of ≥80 Gy.

**Note:** For more information regarding the categories and definitions used for the NCCN Evidence Blocks™, see page [EB-1](#).

All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**Staging—Uterine Carcinomas and Carcinosarcoma****Table 1**

AJCC Tumor-Node-Metastases (TNM) and International Federation of Gynecology and Obstetrics (FIGO) Surgical Staging Systems for Endometrial Cancer

**Definitions for T, N, M**

<b>T</b>	<b>FIGO Stage</b>	<b>Primary Tumor</b>
<b>TX</b>		Primary tumor cannot be assessed
<b>T0</b>		No evidence of primary tumor
<b>Tis</b>		Carcinoma in situ (preinvasive carcinoma)
<b>T1</b>	<b>I</b>	Tumor confined to the corpus uteri, including endocervical glandular involvement
T1a	<b>IA</b>	Tumor limited to the endometrium or invading less than half the myometrium
T1b	<b>IB</b>	Tumor invading one half or more of the myometrium
<b>T2</b>	<b>II</b>	Tumor invading the stromal connective tissue of the cervix but not extending beyond the uterus. Does NOT include endocervical glandular involvement.
<b>T3</b>	<b>III</b>	Tumor involving serosa, adnexa, vagina, or parametrium
T3a	<b>IIIA</b>	Tumor involving the serosa and/or adnexa (direct extension or metastasis)
T3b	<b>IIIB</b>	Vaginal involvement (direct extension or metastasis) or parametrial involvement
<b>T4</b>	<b>IVA</b>	Tumor invading the bladder mucosa and/or bowel mucosa (bullous edema is not sufficient to classify a tumor as T4)

[Continued](#)

Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2017) published by Springer International Publishing.

**Table 1 - Continued**

<b>N</b>	<b>FIGO Stage</b>	<b>Regional Lymph Nodes</b>
<b>NX</b>		Regional lymph nodes cannot be assessed
<b>N0</b>		No regional lymph node metastasis
<b>N0(i+)</b>		Isolated tumor cells in regional lymph node(s) no greater than 0.2 mm
<b>N1</b>	<b>IIIC1</b>	Regional lymph node metastasis to pelvic lymph nodes
N1mi	<b>IIIC1</b>	Regional lymph node metastasis (greater than 0.2 mm but not greater than 2.0 mm in diameter) to pelvic lymph nodes
N1a	<b>IIIC1</b>	Regional lymph node metastasis (greater than 2.0 mm in diameter) to pelvic lymph nodes
<b>N2</b>	<b>IIIC2</b>	Regional lymph node metastasis to para-aortic lymph nodes, with or without positive pelvic lymph nodes
N2mi	<b>IIIC2</b>	Regional lymph node metastasis (greater than 0.2 mm but not greater than 2.0 mm in diameter) to para-aortic lymph nodes, with or without positive pelvic lymph nodes
N2a	<b>IIIC2</b>	Regional lymph node metastasis (greater than 2.0 mm in diameter) to para-aortic lymph nodes, with or without positive pelvic lymph nodes

Suffix (sn) is added to the N category when metastasis is identified only by sentinel lymph node biopsy.

**M FIGO Stage Distant Metastasis**

**M0** No distant metastasis

**M1** Distant metastasis (includes metastasis to inguinal lymph nodes, intraperitoneal disease, lung, liver, or bone). (It excludes metastasis to pelvic or para-aortic lymph nodes, vagina, uterine serosa, or adnexa).

**G Histologic Grade**

**GX** Grade cannot be assessed

**G1** Well differentiated

**G2** Moderately differentiated

**G3** Poorly differentiated or undifferentiated

**Table 2. AJCC Prognostic Stage Groups**

	<b>T</b>	<b>N</b>	<b>M</b>
<b>Stage I</b>	T1	N0	M0
<b>Stage IA</b>	T1a	N0	M0
<b>Stage IB</b>	T1b	N0	M0
<b>Stage II</b>	T2	N0	M0
<b>Stage III</b>	T3	N0	M0
<b>Stage IIIA</b>	T3a	N0	M0
<b>Stage IIIB</b>	T3b	N0	M0
<b>Stage IIIC1</b>	T1-T3	N1/N1mi/N1a	M0
<b>Stage IIIC2</b>	T1-T3	N2/N2mi/N2a	M0
<b>Stage IVA</b>	T4	Any N	M0
<b>Stage IVB</b>	Any T	Any N	M1

[Continued](#)

Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2017) published by Springer International Publishing.

## Staging—Uterine Sarcoma

**Table 3**

AJCC Tumor-Node-Metastases (TNM) and International Federation of Gynecology and Obstetrics (FIGO) Surgical Staging Systems for Uterine Sarcomas (includes Leiomyosarcoma and Endometrial Stromal Sarcoma)

### Leiomyosarcoma and Endometrial Stromal Sarcoma

T	FIGO Stage	Primary Tumor
<b>TX</b>		Primary tumor cannot be assessed
<b>T0</b>		No evidence of primary tumor
<b>T1</b>	<b>I</b>	Tumor limited to the uterus
T1a	<b>IA</b>	Tumor 5 cm or less in greatest dimension
T1b	<b>IB</b>	Tumor more than 5 cm
<b>T2</b>	<b>II</b>	Tumor extends beyond the uterus, within the pelvis
T2a	<b>IIA</b>	Tumor involves adnexa
T2b	<b>IIB</b>	Tumor involves other pelvic tissues
<b>T3</b>	<b>III</b>	Tumor infiltrates abdominal tissues
T3a	<b>IIIA</b>	One site
T3b	<b>IIIB</b>	More than one site
<b>T4</b>	<b>IVA</b>	Tumor invades bladder or rectum

N	FIGO Stage	Regional Lymph Nodes
<b>NX</b>		Regional lymph nodes cannot be assessed
<b>N0</b>		No regional lymph node metastasis
<b>N0(i+)</b>		Isolated tumor cells in regional lymph node(s) no greater than 0.2 mm
<b>N1</b>	<b>IIIC</b>	Regional lymph node metastasis

M	FIGO Stage	Distant Metastasis
<b>M0</b>		No distant metastasis
<b>M1</b>	<b>IVB</b>	Distant metastasis (excluding adnexa, pelvic, and abdominal tissues)

### G Histologic Grade

<b>GX</b>	Grade cannot be assessed
<b>G1</b>	Well differentiated
<b>G2</b>	Moderately differentiated
<b>G3</b>	Poorly differentiated or undifferentiated

**Table 4. AJCC Prognostic Groups**

	T	N	M
<b>Stage I</b>	T1	N0	M0
<b>Stage IA</b>	T1a	N0	M0
<b>Stage IB</b>	T1b	N0	M0
<b>Stage IC</b>	T1c	N0	M0
<b>Stage II</b>	T2	N0	M0
<b>Stage IIIA</b>	T3a	N0	M0
<b>Stage IIIB</b>	T3b	N0	M0
<b>Stage IIIC</b>	T1-3	N1	M0
<b>Stage IVA</b>	T4	Any N	M0
<b>Stage IVB</b>	Any T	Any N	M1

Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2017) published by Springer International Publishing.

### Discussion

This discussion is being updated to correspond with the newly updated algorithm. Partial update 02/08/19, last full update 5/25/18.

#### NCCN Categories of Evidence and Consensus

**Category 1:** Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

**Category 2A:** Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

**Category 2B:** Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

**Category 3:** Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

**All recommendations are category 2A unless otherwise indicated.**

#### NCCN Categories of Preference

**Preferred intervention:** Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability

**Other recommended intervention:** Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes

**Useful in certain circumstances:** Other interventions that may be used for selected patient populations (defined with recommendation)

**All recommendations are considered appropriate.**




# NCCN Guidelines Version 4.2019

## Uterine Neoplasms

[NCCN Guidelines Index](#)  
[Table of Contents](#)  
[Discussion](#)

### Table of Contents

<b>Overview</b> .....	<b>MS-3</b>	Advanced Stage/Extrauterine Disease .....	MS-15
<b>Endometrial Cancer</b> .....	<b>MS-4</b>	Radiotherapy Principles.....	MS-17
Diagnosis and Workup.....	MS-4	Post-Treatment Surveillance .....	MS-17
Disease Staging.....	MS-5	Hormone Replacement Therapy for Hypoestrogenism .....	MS-18
Principles of Evaluation and Surgical Staging for Endometrial Carcinoma.....	MS-5	Treatment of Recurrent or Metastatic Disease .....	MS-19
Pathology.....	MS-6	Localized Disease .....	MS-19
Lymphadenectomy.....	MS-6	Systemic Disease.....	MS-19
Sentinel Lymph Node Mapping .....	MS-7	Hormonal Therapy.....	MS-19
SLN Ultrastaging .....	MS-8	Systemic Therapy.....	MS-20
Minimally Invasive Procedures.....	MS-9	<b>Uterine Serous Carcinomas, Clear Cell Carcinomas, and Carcinosarcomas</b> .....	<b>MS-22</b>
Primary Treatment.....	MS-10	Overview .....	MS-22
Disease Limited to the Uterus .....	MS-10	Treatment.....	MS-22
Incomplete Surgical Staging.....	MS-11	<b>Uterine Sarcomas</b> .....	<b>MS-24</b>
Fertility-Sparing Therapy.....	MS-11	Overview .....	MS-24
Suspected or Gross Cervical Involvement .....	MS-11	Staging and Treatment.....	MS-24
Patients Not Suited for Primary Surgery.....	MS-12	Low-Grade Endometrial Stromal Sarcoma.....	MS-25
Suspected Extrauterine Disease.....	MS-12	High-Grade Endometrial Stromal Sarcoma, Leiomyosarcoma, and Undifferentiated Uterine Sarcoma.....	MS-25
Adjuvant Therapy.....	MS-13	Post-Treatment Surveillance .....	MS-27
Uterine-Confined Disease .....	MS-13	Treatment of Recurrent or Metastatic Disease .....	MS-27
Adjuvant RT .....	MS-14	<b>Drug Reactions</b> .....	<b>MS-28</b>
Adjuvant Systemic Therapy.....	MS-15	<b>References</b> .....	<b>MS-29</b>



### Overview

Adenocarcinoma of the endometrium (also known as endometrial cancer, or more broadly as uterine cancer or carcinoma of the uterine corpus) is the most common malignancy of the female genital tract in the United States. It is estimated that 63,230 new uterine cancer cases will occur in 2018, with 11,350 deaths resulting from the disease.<sup>1</sup> Stromal or mesenchymal sarcomas are uncommon subtypes accounting for approximately 3% of all uterine cancers.<sup>2,3</sup> The NCCN Guidelines for Uterine Neoplasms describe malignant epithelial tumors and uterine sarcomas; each of these major categories contains specific histologic groups that require different management (see *Initial Clinical Findings* in the NCCN Guidelines for Uterine Neoplasms).

Risk factors for uterine neoplasms include increased levels of estrogen (caused by obesity, diabetes, and high-fat diet), early age at menarche, nulliparity, late age at menopause, Lynch syndrome, older age ( $\geq 55$  years), and tamoxifen use.<sup>4-7</sup> Thus, the incidence of endometrial cancer is increasing because of increased life expectancy and obesity. The *Summary of the Guidelines Updates* describes the most recent revisions to the algorithms, which have been incorporated into this revised Discussion text (see the NCCN Guidelines for Uterine Neoplasms). By definition, the NCCN Guidelines cannot incorporate all possible clinical variations and are not intended to replace good clinical judgment or individualization of treatments. Exceptions to the rule were discussed among the NCCN panel during the process of developing these guidelines.

For patients with known or suspected uterine neoplasms, the initial preoperative evaluation/workup for known or suspected malignancy includes a history and physical examination, expert pathology review with additional endometrial biopsy as indicated, imaging, consideration of genetic evaluation, and other studies (see *Initial Evaluation* and *Principles*

*of Imaging* in the NCCN Guidelines for Uterine Neoplasms).<sup>8</sup> Preoperative imaging and biopsy may help to identify uterine sarcomas, although biopsy sensitivity is less than that for endometrial cancer. An expert pathology review will determine whether a patient has a malignant epithelial tumor or a stromal/malignant mesenchymal tumor. Epithelial tumor types include pure endometrioid cancer, uterine serous carcinoma, clear cell carcinoma, carcinosarcoma (also known as malignant mixed Müllerian tumor [MMMT]), and undifferentiated/dedifferentiated carcinoma. Stromal or mesenchymal tumor types (interchangeable terms) include uterine leiomyosarcoma (uLMS), endometrial stromal sarcoma (ESS), undifferentiated uterine sarcoma (UUS, previously called high-grade undifferentiated endometrial sarcoma), adenosarcoma, and perivascular epithelioid cell neoplasm (PEComa). Given the typical age group at risk for uterine neoplasms (ie,  $\geq 55$  years) and the presence of comorbid illnesses in older patients, it is prudent in selected patients to also measure renal and liver function.

Most endometrial cancer is caused by sporadic mutations. However, genetic mutations cause endometrial cancer in about 5% of patients, which occurs 10 to 20 years before sporadic cancer.<sup>9</sup> Screening for genetic mutations (eg, Lynch syndrome/hereditary non-polyposis colorectal cancer) should be considered in all patients with endometrial (and colorectal) cancer but especially in those younger than 50 years of age.<sup>7,9-11</sup> Genetic testing and counseling should be considered for patients younger than 50 years of age with endometrial cancer and those with a significant family history of endometrial and/or colorectal cancer.<sup>12-14</sup> If these patients have Lynch syndrome, they are at greater risk for a second cancer (eg, colorectal cancer, ovarian cancer).<sup>5,11,15</sup> In addition, their relatives may have Lynch syndrome.

Screening of the tumor for defective DNA mismatch repair (MMR) using immunohistochemistry and/or microsatellite instability (MSI) is used to

identify which patients should undergo mutation testing for Lynch syndrome (see *Lynch Syndrome* in the NCCN Guidelines for Colorectal Cancer Screening).<sup>9,10,16-20</sup> Universal testing of endometrial tumors for defects in DNA MMR is recommended (eg, *MLH1*, *MSH2*, *MSH6*). *MLH1* loss should be further evaluated for promoter methylation to assess for an epigenetic process rather than a germline mutation.<sup>18</sup> Genetic counseling and testing is recommended for patients with all other MMR abnormalities and for patients without MMR defects but who have a significant family history of endometrial and/or colorectal cancer (See *Lynch Syndrome [Hereditary Non-Polyposis Colorectal Cancer]* in the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal).

Women with Lynch syndrome are at higher lifetime risk (up to 60%) for endometrial cancer; thus, close monitoring and discussion of risk-reducing strategies is recommended.<sup>10,21,22</sup> In relatives with Lynch syndrome but without endometrial cancer, a yearly endometrial biopsy is recommended to assess for cancer.<sup>13,23</sup> This strategy also enables select women to defer surgery (and surgical menopause) and to preserve their fertility. Prophylactic hysterectomy/bilateral salpingo-oophorectomy (BSO) can then be done after childbearing is complete or sooner, depending on patient preference.<sup>24,25</sup> In addition, interventions to decrease the risk from colorectal cancer may also be appropriate (eg, annual colonoscopy).

### Endometrial Cancer

In 2017, 67% of patients with adenocarcinoma of the endometrium were diagnosed with disease confined to the uterus at diagnosis.<sup>26</sup> Regional and distant disease comprised 21% and 8% of cases, respectively.

Many physicians believe that adenocarcinoma of the endometrium is a more treatable malignancy because the early symptoms of irregular vaginal bleeding (in this predominantly postmenopausal patient population) often trigger patients to seek care when the disease is at an

early and treatable stage. However, data show that the mortality rate for uterine cancer has increased more rapidly than the incidence rate.<sup>27</sup> This increased mortality may be related to an increased rate of advanced-stage cancers, high-risk histologies (eg, serous carcinomas), and patients being diagnosed at an older age. Analysis of SEER data suggests that survival is increased in patients who are younger, have early-stage disease, and have lower-grade disease.<sup>28</sup> In addition to grade and depth of myometrial invasion, other risk factors associated with poor prognosis include age, lymph node status, tumor size, lymphovascular space invasion (LVSI), and tumor involvement of the lower uterine segment.<sup>29,30</sup> To further improve outcome for patients with this disease, physicians need to identify high-risk patients and to tailor treatment appropriately to provide the best long-term survival. The panel suggests that gynecologic oncologists be involved in the primary management of all patients with endometrial cancer.

### Diagnosis and Workup

About 90% of patients with endometrial carcinoma have abnormal vaginal bleeding, most commonly in the postmenopausal period. The workup was previously described (see *Overview* in this Discussion). Diagnosis can usually be made by an office endometrial biopsy.<sup>31,32</sup> The histologic information from the endometrial biopsy (with or without endocervical curettage) should be sufficient for planning definitive treatment. Office endometrial biopsies have a false-negative rate of about 10%. Thus, a negative endometrial biopsy in a symptomatic patient must be followed by a fractional dilation and curettage (D&C) under anesthesia.<sup>31,33</sup> Hysteroscopy may be helpful in evaluating the endometrium for lesions, such as a polyp, if the patient has persistent or recurrent undiagnosed bleeding.<sup>34</sup> Endometrial biopsy may not be accurate for diagnosing malignancies of the uterine wall such as mesenchymal tumors.

For detailed imaging recommendations by stage and planned treatment approach, see *Principles of Imaging* in the NCCN Guidelines for Uterine

Neoplasms. Consideration of chest imaging (chest x-ray) is recommended. Other imaging tests such as CT, MRI, and/or PET/CT may be used to assess disease extent and to evaluate for metastatic disease as indicated based on clinical symptoms, physical findings, or abnormal laboratory findings.<sup>35-40</sup> In patients with extrauterine disease, a serum CA-125 assay may be helpful in monitoring clinical response.<sup>41,42</sup> However, serum CA-125 levels can be falsely increased in women who have peritoneal inflammation/infection or radiation injury, may be normal in women with isolated vaginal metastases, and may not predict recurrence in the absence of other clinical findings.<sup>43-45</sup> Currently, there is no validated screening test for endometrial carcinoma.<sup>46,47</sup>

### Disease Staging

The FIGO (International Federation of Gynecology and Obstetrics) system is most commonly used for staging uterine cancer. The original 1970 criteria for staging endometrial cancer only used information gained from presurgical evaluation (including physical examination and diagnostic fractional D&C). At that time, many patients were not treated with primary surgery because of obesity or various other medical problems. Thus, the 1970 staging system is rarely used today (eg, when the patient is not a surgical candidate).

Several studies demonstrated that clinical staging was inaccurate and did not reflect actual disease extent in 15% to 20% of patients.<sup>48-50</sup> This reported understaging and, more importantly, the ability to identify multiple prognostic factors with a full pathologic review made possible with surgical staging, motivated a change in the staging classification. Therefore, in 1988, FIGO modified its staging system to emphasize thorough surgical/pathologic assessment of data, such as histologic grade, myometrial invasion, and the extent and location of extrauterine spread (including retroperitoneal lymph node metastases).<sup>51</sup> FIGO updated and refined the surgical/pathologic staging criteria for uterine neoplasms in

2009.<sup>52-55</sup> Separate staging systems for malignant epithelial tumors and uterine sarcomas are now available (see *Staging* section of the algorithm). In 2017, the AJCC Cancer Staging Manual was updated (to take effect January 2018).<sup>56</sup>

The 2009 FIGO staging system streamlined stages I and II endometrial carcinoma. These revisions were made because the survival rates for some of the previous sub-stages were similar.<sup>54</sup> Stage IA is now less than 50% myometrial invasion, and stage IB is 50% or more myometrial invasion. Stage II only includes patients with cervical stromal invasion. Patients with uterine-confined disease and endocervical glandular involvement (mucosal involvement) without cervical stromal invasion are no longer considered stage II.<sup>54</sup> Stage IIIC is now subdivided into IIIC1 and IIIC2, because survival is worse with positive para-aortic nodes.<sup>54</sup> While most of the previously published studies discussed in these NCCN Guidelines used the older 1988 FIGO staging system, these have been reinterpreted by the NCCN panel to reconcile with the 2009 staging system.

Peritoneal cytology no longer affects the 2009 FIGO staging, because it is not viewed by some authors as an independent risk factor.<sup>55</sup> However, FIGO and AJCC continue to recommend that peritoneal washings be obtained and results recorded, because positive cytology may add to the effect of other risk factors (see *Principles of Evaluation and Surgical Staging* in the NCCN Guidelines for Endometrial Carcinoma).<sup>57,58</sup>

### Principles of Evaluation and Surgical Staging for Endometrial Carcinoma

Staging should be done by a team with expertise in imaging, pathology, and surgery. The amount of surgical staging that is necessary to determine disease status depends on preoperative and intraoperative assessment of findings by experienced surgeons. For the 2014 update, the NCCN panel added a new section on surgical staging (see *Principles*

of Evaluation and Surgical Staging in the NCCN Guidelines for Endometrial Carcinoma). However, this surgical staging section only applies to malignant epithelial tumors and not to uterine sarcomas. Surgical staging with nodal assessment for apparent uterine-confined endometrial cancer is critical to accurately determine the initial FIGO stage. The NCCN sentinel lymph node (SLN) algorithm is recommended if sentinel node mapping is utilized.

### Pathology

An expert pathology review will determine the specific epithelial histology of the tumor (ie, various endometrioid histologies, serous carcinoma, clear cell carcinoma, carcinosarcoma, undifferentiated carcinoma). The pathologic assessment of the uterus and the nodes is described in the algorithm; this assessment should also include the Fallopian tubes, ovaries, and peritoneal cytology. If nodal resection was performed, the level of nodal involvement and size of metastasis should be determined. See *Hysterectomy and Pathologic Evaluation* in the NCCN Guidelines for Endometrial Carcinoma. The *Protocol for Examination of Specimens from Patients With Carcinoma and Carcinosarcoma of the Endometrium* from the College of American Pathologists (CAP) is a useful guide (<https://cap.objects.frb.io/protocols/cp-femalereproductive-endometrium-18protocol-4100.pdf>). Estrogen receptor testing is recommended in the setting of stage III, IV, or recurrent endometrioid carcinoma.

As the grade of the tumor increases, the accuracy of intraoperative evaluation of myometrial invasion decreases (ie, assessment by gross examination of fresh tissue). In one study, the depth of invasion was accurately determined by gross examinations in 87.3% of grade 1 lesions, 64.9% of grade 2 lesions, and 30.8% of grade 3 lesions.<sup>59</sup> Studies show that in 15% to 20% of cases, the preoperative grade (as assessed by endometrial biopsy or curettage) is upgraded on final fixed pathologic evaluation of the hysterectomy specimen.<sup>60</sup>

### Lymphadenectomy

Previously, a full standard lymphadenectomy (ie, dissection and assessment of both pelvic and para-aortic nodes) was recommended for all patients; however, a more selective and tailored lymphadenectomy approach that may include the SLN algorithm is now recommended by the NCCN Panel to avoid systematic overtreatment.<sup>61</sup> No randomized trial data support routine full lymphadenectomy,<sup>62</sup> although some retrospective studies have suggested that it is beneficial.<sup>63-65</sup> Two randomized clinical trials from Europe reported that routine lymph node dissection did not improve the outcome of endometrial cancer patients, but lymphadenectomy did identify those with nodal disease.<sup>66,67</sup> However, these findings remain a point of contention.<sup>68-70</sup> To avoid over-interpretation of these results, it is important to address the limitations of these randomized studies, including selection of patients, extent of lymph node dissection, and standardization of postoperative therapy.<sup>71,72</sup> The other concerns include the lack of central pathology review, subspecialty of surgeons, and adequacy of statistical power.

Decisions about whether to perform lymphadenectomy, and, if done, to what extent (eg, pelvic nodes only or both pelvic and para-aortic nodes), can be made based on preoperative and intraoperative findings. Criteria have been suggested as indicative of low risk for nodal metastases: 1) less than 50% myometrial invasion; 2) tumor less than 2 cm; and 3) well or moderately differentiated histology.<sup>73,74</sup> However, this may be difficult to accurately determine before final pathology results are available.

Another associated benefit of lymphadenectomy is the diagnosis of those with nodal metastases to guide appropriate adjuvant treatment to improve survival or decrease toxicity. However, one of the trials was not designed to address this question.<sup>67</sup> Therefore, there was no standardization of adjuvant treatment after staging surgery with lymphadenectomy. In fact, the use of lymphadenectomy did not translate into an increased use of

adjuvant therapy. This may have contributed to the lack of difference in recurrence and survival in the two groups.

The question of whether to add para-aortic lymphadenectomy to pelvic node dissection has been debated. Prior studies have shown conflicting information regarding the risk of para-aortic nodal metastases in patients without disease in the pelvic nodes.<sup>50,73,75,76</sup> There was a high rate of lymphatic metastasis above the inferior mesenteric artery, suggesting a need for systematic pelvic and para-aortic lymphadenectomy. Hence, para-aortic lymphadenectomy up to the renal vessels may be considered for selective high-risk situations, including those with pelvic lymphadenectomy or high-risk histologic features. Many surgeons do not do a full lymphadenectomy in patients with grade 1 early-stage endometrial cancer.<sup>61</sup>

In summary, lymph node dissection identifies patients requiring adjuvant treatment with RT and/or systemic therapy.<sup>77</sup> A subset of patients may not benefit from lymphadenectomy; however, it is difficult to preoperatively identify these patients because of the uncontrollable variables of change in grade and depth of invasion on final pathology. The NCCN panel recommends that lymphadenectomy should be done for selected patients with endometrial cancer with para-aortic lymphadenectomy done as indicated for high-risk patients (see *Principles of Evaluation and Surgical Staging* in the NCCN Guidelines for Endometrial Carcinoma).<sup>6</sup> Lymphadenectomy is contraindicated for patients with uterine sarcoma. SLN mapping can be considered as an alternative to full lymphadenectomy in the setting of apparent uterine-confined disease. The SLN surgical algorithm is described below.

### *Sentinel Lymph Node Mapping*

The section on surgical staging (see *Principles of Evaluation and Surgical Staging* in the NCCN Guidelines for Endometrial Carcinoma) includes recommendations about SLN mapping. SLN mapping may be considered

for patients with apparent uterine-confined endometrial cancer to assess whether they have metastatic pelvic lymph nodes.<sup>78-82</sup> In SLN mapping, dye is injected into the cervix, which travels to the sentinel nodes (see Figures 1–3 in *Principles of Evaluation and Surgical Staging* in the NCCN Guidelines for Endometrial Carcinoma).

A surgical SLN algorithm is proposed to decrease the false-negative rate (see Figure 4 in *Principles of Evaluation and Surgical Staging* in the NCCN Guidelines for Endometrial Carcinoma).<sup>78,83</sup> For example, suspicious or grossly enlarged nodes should be removed regardless of SLN mapping results. In SLN mapping, the surgeon's expertise and attention to technical detail are critical. Patients may be able to avoid the morbidity of a standard lymphadenectomy with SLN mapping.<sup>84,85</sup> Because SLNs identify the primary lymphatic pathway, this increases the yield of finding metastatic disease during the mapping process. If SLN mapping fails, a reflex side-specific nodal dissection should be performed.<sup>78,86</sup> SLN mapping may be most appropriate for those at low to intermediate risk for metastases and/or for those who may not tolerate a standard lymphadenectomy.<sup>82,85-91</sup> Recent findings also suggest that indocyanine green may be preferable to blue dyes.<sup>91-95</sup> Attention to detail and experience are critical to ensure optimal outcomes.

An updated literature review and consensus recommendations for SLN mapping in endometrial cancer were recently released by the Society of Gynecologic Oncology (SGO).<sup>82</sup> Close adherence to the NCCN SLN surgical algorithm was found to result in accurate prediction of pelvic lymph node metastasis with a less than 5% false-negative rate. Additionally, results were recently published from the FIRES trial, which compared SLN mapping to lymphadenectomy for endometrial cancer in the largest multicenter prospective study to date (n = 385).<sup>91</sup> Mapping of at least one SLN was successful in 86% of patients; sensitivity was 97.2%

(95% CI, 85.0–100), and negative predictive value was 99.6% (95% CI, 97.9–100).

Until recently, much of the data to support SLN mapping was based on single-institution studies. A systematic review of 17 studies with  $n > 30$  patients revealed detection rates of 60% to 100%; detection rates for studies with larger cohorts ( $n > 100$ ) were at least 80%. Retrospective application of a surgical algorithm generated 95% sensitivity, 99% predictive value, and a 5% false-negative rate.<sup>96</sup> Another recent systematic review and meta-analysis of 55 studies with  $n > 10$  patients ( $n = 4915$ ) generated an overall detection rate of 81% with a 50% bilateral pelvic node detection rate and 17% paraaortic detection rate.<sup>95</sup>

SLN mapping should be done in institutions with expertise in this procedure. If patients have apparent metastatic disease (based on imaging and surgical exploration), removal of nodes for staging purposes is not necessary because it will not change management.<sup>35</sup> The main contraindication for SLN mapping is uterine sarcoma. Historically, SLN mapping was controversial in patients with high-risk histology (eg, serous carcinoma, clear cell carcinoma, carcinosarcoma).<sup>61,97</sup> However, recently, SLN mapping in patients with high-risk histologies (ie, grade 3, serous, clear cell, carcinosarcoma) has been reported with promising results as a potential alternative to complete lymphadenectomy.<sup>86,98</sup>

### **SLN Ultrastaging**

Recent data highlight the potential significance and impact of SLN ultrastaging (ie, serial sectioning and immunohistochemistry) to improve the accuracy of detecting micrometastases. Ultrastaging of SLNs can reveal lymph node metastases undetected through conventional histology, and studies suggest that SLN ultrastaging leads to upstaging in 5% to 15% of patients.<sup>81,84,88,90,96</sup>

In a retrospective analysis of patients with early-stage endometrial cancer ( $n = 780$ ) who underwent SLN mapping with lymphadenectomy versus lymphadenectomy alone, SLN mapping led to the detection of more metastasis (30.3% vs. 14.7%,  $P < .001$ ) and was associated with greater use of adjuvant therapy.<sup>99</sup> Long-term follow-up was reported from a prospective multicenter study in 125 patients with early-stage endometrial carcinoma who underwent SLN biopsy. Patients with a positive SLN underwent external beam radiation therapy (EBRT) and chemotherapy at a higher rate than those with a negative SLN. In patients with a detected SLN, recurrence-free survival at 50 months was 84.7%, and no difference was detected between patients with and without a positive SLN ( $P = .5$ ).<sup>100</sup>

In a cohort of 508 patients who underwent SLN mapping, ultrastaging detected 23 additional cases of micrometastasis that would have been missed by conventional hematoxylin and eosin staining.<sup>101</sup> A multicenter study of 304 women with presumed low- or intermediate-risk disease showed that SLN biopsy and ultrastaging detected metastatic SLNs in a 3-fold greater number of patients than standard lymphadenectomy.<sup>102</sup>

Although these findings do not appear to be an artifact of uterine manipulation,<sup>103</sup> the implications and appropriate management of micrometastases or isolated tumor cells (ITCs) detected via SLN ultrastaging are not yet clear.<sup>82,84,90,104-106</sup> The prognostic significance of ITCs has been studied in breast cancer<sup>107</sup> where nodes containing ITCs are excluded from the positive node count per AJCC staging. Studies have recently begun to investigate the significance of ITCs discovered during SLN mapping in early-stage endometrial cancer.

A retrospective review examined 844 patients with endometrial cancer that underwent SLN mapping.<sup>108</sup> The majority of patients with ITCs, micrometastasis, and macrometastasis received adjuvant chemotherapy (83%, 81%, and 89%, respectively). Recurrence-free survival at 3 years was 90% for those with negative SLNs, 86% for ITCs, and 86% for

micrometastasis. Only patients with SLN macrometastasis had significantly lower recurrence-free survival (71%,  $P < .001$ ).

A recent prospective observational study of 519 patients compared outcomes for patients with SLN macrometastasis, micrometastasis, and ITCs, taking into account adjuvant treatment.<sup>109</sup> Patients with SLN ITCs had a significantly better 3-year progression-free survival (PFS) compared with patients with SLN macrometastasis (95.5% vs. 58.5%), and outcomes were similar between patients with negative SLNs, ITCs, and micrometastasis. Recurrence was detected in only 1 of 31 patients with ITCs (stage IB carcinosarcoma) and adjuvant treatment did not appear to influence outcomes. Based on these early data, it is unclear if patients with SLN ITCs would derive significant benefit from adjuvant treatment. Future evaluation of prognosis/outcome may need to prospectively examine the threshold for and impact of adjuvant therapy for patients with scattered ITCs.

### **Minimally Invasive Procedures**

Over the past decade, practice has trended towards minimally invasive approaches to total hysterectomy (TH)/BSO and lymph node assessment in patients with early-stage endometrial cancer.<sup>110</sup> Although these procedures may be performed by any surgical route (eg, laparoscopic, robotic, vaginal, abdominal), the standard in those with apparent uterine-confined disease is to perform the procedure via a minimally invasive approach. Randomized trials, a Cochrane Database Systematic Review, and population-based surgical studies support that minimally invasive techniques are preferred in this setting due to a lower rate of surgical site infection, transfusion, venous thromboembolism, decreased hospital stay, and lower cost of care, without compromise in oncologic outcome.<sup>110-116</sup> Despite data showing that minimally invasive procedures result in lower perioperative complications and lower cost of care, racial and geographic

disparities in access to minimally invasive surgical care have been observed.<sup>112,116</sup>

A randomized phase III trial evaluated laparoscopy for comprehensive surgical staging; patients ( $n = 2616$ ) with clinical stage I to IIA disease (GOG-LAP2) were assessed.<sup>115,117</sup> Patients were randomly allocated 2:1 to laparoscopy or laparotomy. Results from LAP2 indicate that 26% of patients needed conversion to laparotomy because of poor visibility, metastatic cancer, bleeding, increased age, or increased body mass index. Detection of advanced cancer was not significantly different between the groups. However, significant differences were noted in removal of pelvic and para-aortic nodes (8% not removed with laparoscopy vs. 4% with laparotomy,  $P < .0001$ ).<sup>118,119</sup> Significantly fewer postoperative adverse events and shorter hospitalization occurred with laparoscopy compared with laparotomy. Recurrence rates were 11.4% for laparoscopy versus 10.2% for laparotomy. The 5-year overall survival (OS) rate was 84.8% for both arms of LAP2.<sup>117</sup> Laparoscopic staging was associated with improved postoperative quality of life across several parameters.<sup>114</sup>

Results were recently published from the LACE trial, which compared outcomes of patients with stage I endometrial carcinoma ( $n = 760$ ) who were randomized to undergo TAH or total laparoscopic hysterectomy.<sup>111</sup> At a median follow-up of 4.5 years, disease-free survival (DFS) was 81.3% for laparotomy versus 81.6% for laparoscopy, with no significant differences observed between groups for recurrence and OS. Another randomized trial ( $n = 283$ ) comparing laparoscopy versus laparotomy reported shorter hospital stay, less pain, and faster resumption of daily activities with laparoscopy.<sup>120</sup> However, laparotomy may still be required for certain clinical situations (eg, elderly patients, those with a very large uterus) or certain metastatic presentations.<sup>115,121,122</sup>

Robotic surgery is a minimally invasive technology that has been increasingly used in the surgical staging of early-stage endometrial carcinoma due to its potential advantages over laparotomy, especially for obese patients.<sup>123-127</sup> Prospective cohort and retrospective studies suggest that robotic approaches perform similarly to laparoscopy and result in comparable or improved perioperative outcomes.<sup>127-130</sup> Oncologic outcomes appear to be comparable to other surgical approaches, although longer-term outcomes are still being investigated.<sup>131-133</sup> In heavier patients, robotic surgery may result in less frequent conversion to laparotomy when compared with laparoscopic approaches and also appears to be safe and feasible in patients at higher anesthesiologic risk.<sup>127,128,134</sup>

Costs for robotic equipment and maintenance remain high.<sup>135-123,124,131-133,136</sup> The SGO, American Association of Gynecologic Laparoscopists (AAGL), and American Congress of Obstetricians and Gynecologists (ACOG) have published guidelines or position statements about robotic surgery.<sup>137-139</sup> For recent reviews on the robotic-assisted surgery for gynecologic malignancies and associated cost issues, see Sinno and Fader and Gala et al.<sup>140,141</sup>

### Primary Treatment

These NCCN Guidelines divide pure endometrioid cancer into three categories for delineating treatment: 1) disease limited to the uterus; 2) suspected or gross cervical involvement; and 3) suspected extrauterine disease. Most patients with endometrial cancer have stage I disease at presentation, and surgery (with or without adjuvant therapy) is recommended for medically operable patients. As a general principle, endometrial carcinoma should be removed en bloc to optimize outcomes; intraperitoneal morcellation should be avoided.<sup>142-145</sup>

### Disease Limited to the Uterus

To stage medically operable patients with endometrioid histologies clinically confined to the fundal portion of the uterus, the recommended surgical procedure includes TH/BSO with surgical staging and lymph node assessment (see *Hysterectomy and Pathologic Evaluation*, and *Principles of Evaluation and Surgical Staging* in the NCCN Guidelines for Endometrial Carcinoma and in this Discussion).<sup>68</sup> When indicated, surgical staging is recommended to gather full pathologic and prognostic data on which to base decisions regarding adjuvant treatment for select patients who do not have medical or technical contraindications to lymph node dissection (see *Lymphadenectomy* and *Sentinel Lymph Node Mapping* in this Discussion). Ovarian preservation may be safe in select premenopausal women with stage I endometrioid cancer.<sup>146-148</sup> Minimally invasive surgery is the preferred approach when technically feasible and is considered a quality measure by the SGO and the American College of Surgeons ([www.sgo.org/quality-outcomes-and-research/quality-indicators](http://www.sgo.org/quality-outcomes-and-research/quality-indicators); [www.facs.org/quality-programs/cancer/ncdb/qualitymeasures](http://www.facs.org/quality-programs/cancer/ncdb/qualitymeasures)).

During surgery, the intraperitoneal structures should be carefully evaluated, and suspicious areas should be biopsied. While not specifically affecting staging, FIGO recommends that peritoneal cytology should be collected and results should be recorded. Enlarged or suspicious lymph nodes should be excised to confirm or rule out metastatic disease. Retroperitoneal node dissection with pathologic evaluation—in the absence of clinically apparent lymphadenectomy—is useful when using the 2009 FIGO staging criteria, but its routine use has been questioned (see *Lymphadenectomy* in this Discussion).

Patients with apparent uterine-confined endometrial carcinoma are candidates for sentinel node mapping, which assesses the pelvic nodes bilaterally and may be less morbid than complete lymphadenectomy (see

*Sentinel Lymph Node Mapping* in this Discussion). Adherence to the NCCN SLN algorithm is critical.

### *Incomplete Surgical Staging*

For patients with incomplete (ie, not thorough) surgical staging and high-risk intrauterine features, imaging is often recommended, especially in patients with higher grade and more deeply invasive tumors.<sup>149,150</sup> Surgical restaging, including lymph node dissection, can also be done.<sup>73</sup> Based on the imaging and/or surgical restaging results, recommended adjuvant treatment options are provided in the algorithm (see Adjuvant Treatment for *Incompletely Surgically Staged* in the NCCN Guidelines for Endometrial Carcinoma).

### *Fertility-Sparing Therapy*

Although the primary treatment of endometrial cancer is usually hysterectomy, continuous progestin-based therapy may be considered for highly selected patients with Grade 1, stage IA (noninvasive) disease who wish to preserve their fertility.<sup>151-155</sup> Likewise, it may also be selectively used for young patients with endometrial hyperplasia who desire fertility preservation. The guidelines include an algorithm for fertility-sparing therapy in selected patients with biopsy-proven grade 1 (preferably by D&C), stage IA noninvasive endometrioid adenocarcinoma (see *Criteria for Considering Fertility-Sparing Options* in the NCCN Guidelines for Endometrial Carcinoma). The panel recommends consultation with a fertility expert. When considering fertility-sparing therapy, all of the criteria must be met as outlined in the algorithm (eg, no metastatic disease). Selected patients may require genetic counseling and testing. Patients should also receive counseling that fertility-sparing therapy is not the standard of care for the treatment of endometrial carcinoma. TH/BSO with surgical staging is recommended after childbearing is complete, if therapy is not effective, or if progression occurs. Fertility-sparing therapy is not recommended for high-risk patients (eg, those with high-grade

endometrioid adenocarcinomas, uterine serous carcinoma, clear cell carcinoma, carcinosarcoma, and uLMS).

Continuous progestin-based therapy may include megestrol acetate, medroxyprogesterone, or an intrauterine device containing levonorgestrel.<sup>151,152,156</sup> A durable complete response occurs in about 50% of patients.<sup>151</sup> The use of progestin-based therapy should be carefully considered in the context of other patient-specific factors, including contraindications such as breast cancer, stroke, myocardial infarction, pulmonary embolism, deep vein thrombosis, and smoking.

In patients receiving progestin-based therapies, the NCCN panel recommends close monitoring with endometrial sampling (biopsies or D&C) every 3 to 6 months. TH/BSO with staging is recommended: 1) after childbearing is complete; 2) if patients have documented progression on the biopsies; or 3) if endometrial cancer is still present after 6 to 12 months of progestin-based therapy.<sup>155,157</sup> Although some young women who had subsequent negative endometrial biopsies after hormonal therapy were able to become pregnant (35%), their ultimate recurrence rate was high (35%).<sup>151,154,158-160</sup> In patients with persistent endometrial carcinoma after 6 months of failed hormonal therapy, the panel recommends pelvic MRI as to exclude myoinvasion and nodal/ovarian metastasis before continuing on fertility sparing therapy.

In premenopausal women with stage IA to B endometrial cancer, data suggest that ovarian preservation is safe and not associated with an increased risk of cancer-related mortality; patients were followed for 16 years.<sup>146</sup> Other studies also suggest that ovarian preservation may be safe in women with early-stage endometrial cancer.<sup>147,148</sup>

### ***Suspected or Gross Cervical Involvement***

For patients with suspected or gross cervical involvement (endometrioid histologies), cervical biopsy or pelvic MRI should be performed if not done

previously (see *Additional Workup* in the NCCN Guidelines for Endometrial Carcinoma).<sup>149,150</sup> If negative, patients are assumed to have disease that is limited to the uterus and are treated as previously described (see *Primary Treatment* in the NCCN Guidelines for Endometrial Carcinoma). It may be difficult to distinguish primary cervical carcinoma from stage II endometrial carcinoma. Thus, for operable patients with cervical involvement, TH or radical hysterectomy is recommended along with BSO, cytology (peritoneal lavage), and dissection of lymph nodes if indicated (see *Principles of Evaluation and Surgical Staging and Hysterectomy and Pathologic Evaluation* in the NCCN Guidelines for Endometrial Carcinoma).<sup>68</sup> In these patients, radical or modified radical hysterectomy may improve local control and survival when compared with TH.<sup>161,162</sup> Alternatively, the patient may undergo EBRT and brachytherapy (category 2B) followed by TH/BSO and surgical staging. However, preoperative RT is a category 2B recommendation because the NCCN panel feels that upfront surgery is the preferred option for these patients.

### *Patients Not Suited for Primary Surgery*

For uterine-confined disease not suitable for primary surgery, EBRT and/or brachytherapy is the preferred treatment approach. Initial systemic therapy can also be considered for selected patients with uterine-confined tumors of endometrioid histology (eg, estrogen and progesterone receptor–positive [ER/PR-positive]). Patients receiving hormonal therapy alone should be closely monitored by endometrial biopsy (eg, consider endometrial biopsies every 3–6 months).<sup>46,163</sup> Progesterone-based therapy has been shown to provide some benefit with low toxicity in patients with low-grade tumors.<sup>164</sup> Tamoxifen with alternating megestrol<sup>165</sup> and aromatase inhibitors have also been used.<sup>166-169</sup>

For suspected gross cervical involvement in patients who are not suited for primary surgery, EBRT and brachytherapy is an effective treatment that

can provide some measure of pelvic control and long-term PFS (see *Principles of Radiation Therapy* in the NCCN Guidelines for Uterine Neoplasms).<sup>170-173</sup> EBRT and brachytherapy should be administered with (or without) systemic therapy. If rendered operable, local treatment should follow. Systemic therapy alone is also a primary treatment option (category 2B), but should be followed by local treatment consisting of surgery if feasible (EBRT + brachytherapy if inoperable).

### **Suspected Extrauterine Disease**

If extrauterine disease (endometrioid histologies) is suspected, imaging studies are recommended if clinically indicated (see *Additional Workup* in the NCCN Guidelines for Endometrial Carcinoma) and CA-125 testing can be considered. Estrogen receptor testing is recommended in the setting of stage III or IV endometrioid tumors. Patients with no evidence of extrauterine disease are treated using the guidelines for disease limited to the uterus. Ascites or disease with involvement of the omentum, nodes (including inguinal nodes), ovaries, or peritoneum warrants surgical intervention using TH/BSO with cytology (peritoneal lavage), pelvic and para-aortic lymph node dissection if indicated, and surgical debulking. Consider preoperative systemic therapy. The surgical goal is to have no measurable residual disease; several studies support debulking.<sup>68,174-176</sup>

Patients with unresectable extrauterine pelvic disease (ie, vaginal, bladder, bowel/rectal, nodal, or parametrial involvement) are typically treated with EBRT with (or without) brachytherapy and/or systemic therapy, followed by re-evaluation of tailored surgery.<sup>177-180</sup> Systemic therapy alone can also be considered. Based on treatment response, patients should be re-evaluated for surgical resection and/or RT. For distant visceral metastasis (eg, liver involvement), recommended options include systemic therapy and/or EBRT and/or hormone therapy. Palliative TH/BSO may be considered.

### Adjuvant Therapy

#### ***Uterine-Confined Disease***

Thorough surgical staging provides important information to assist in selection of adjuvant therapy for endometrial tumors (see *Principles of Evaluation and Surgical Staging* in the NCCN Guidelines for Endometrial Carcinoma). Patients with stage I endometrial cancer who have thorough surgical staging are stratified by adverse risk factors (ie, age, positive LVSI, tumor size, and lower uterine segment or surface glandular involvement).<sup>181,182</sup> Recommended adjuvant treatment is shown in the algorithm (see the NCCN Guidelines for Endometrial Carcinoma). Note that the treatment algorithm was revised in 2010 based on the updated FIGO staging.<sup>54</sup> However, by necessity, much of the discussion in this manuscript has been based on data from patients staged using the older FIGO/AJCC staging system. The implications of *stage migration* should be taken into account when evaluating historical data.

The basic concept underlying the recommendations in the NCCN Guidelines is the trend toward selection of more aggressive adjuvant therapy for patients as tumor grade and myometrial and/or cervical invasion worsen, because risk exists on a continuum.<sup>183-185</sup> In surgical stage I and II endometrial cancer, other pathologic factors that may influence the decision regarding adjuvant therapy include LVSI, patient age, tumor volume, depth of invasion, and lower uterine segment or surface cervical glandular involvement. When administering adjuvant RT, it should be initiated as soon as the vaginal cuff has healed, no later than 12 weeks after surgery.

Significant controversy centers on how much adjuvant therapy is necessary in patients with surgical stage I endometrial cancer, regardless of intrauterine features, if extrauterine disease has been clearly ruled out. In a large prospective study, the Gynecologic Oncology Group (GOG) reported that the 5-year survival rate for surgical stage I patients with no

adverse risk factors other than grade and myometrial invasion (ie, without extrauterine disease, isthmus/cervical involvement, or LVSI) was 92.7%.<sup>186</sup>

The practice of surgical staging has led to a decrease in the use of adjuvant therapy for stage I endometrial carcinoma, which is reflected in the option of *observation* in the NCCN Guidelines (see section on adjuvant treatment in the NCCN Guidelines for Endometrial Carcinoma).<sup>77,182,183,187-</sup>

<sup>189</sup> The NCCN panel recommends observation only for select patients with no residual disease in the hysterectomy specimen.

The recommended postoperative (ie, adjuvant) treatment options for surgical stage II patients (using thorough surgical staging) are shown in the algorithm (see *Adjuvant Treatment for Stage II* in the NCCN Guidelines for Endometrial Carcinoma). The NCCN panel generally agrees on the role of adjuvant therapy for patients with an invasive cervical component if extrafascial hysterectomy is performed. However, for patients with stage II disease who have had a radical hysterectomy with negative surgical margins and no evidence of extrauterine disease, observation or vaginal brachytherapy are options. As with stage I disease, the presence of adverse risk factors should be considered when selecting adjuvant therapy.<sup>190</sup>

In 2017, the panel removed observation as a recommended option in the adjuvant setting for patients with stage IA, grade 3 disease with additional risk factors and stage IB grade 3 disease without adverse risk factors. For patients with stage IA/IB, grade 3 disease (IA with adverse risk factors and IB without), systemic therapy was added as a category 2B option when performed along with the primary recommendation of vaginal brachytherapy and/or EBRT. For stage IB, grade 3 disease with adverse risk factors, the option of systemic therapy (in addition to EBRT and/or vaginal brachytherapy) was upgraded to a category 2A option.

### Adjuvant RT

Several phase III trials have assessed adjuvant therapy in patients with uterine-confined disease. In summary, the use of adjuvant RT improves pelvic control in patients with selected risk factors (and may improve PFS), but RT did not improve OS in any of the trials. However, many of these trials had limitations because most of the patients were low risk (ie, they had low-risk intrauterine pathologic risk factors). Thus, the trials were underpowered for patients with high-risk factors. It is recognized that in patients with uterine-confined disease, there is a spectrum of risk based on intrauterine pathologic findings. Adverse intrauterine pathologic risk factors include high-grade tumors, deep myometrial invasion (and consequently more advanced stage), LVSI, and serous or clear cell carcinoma histologies.

Four trials have evaluated the role of adjuvant external-beam pelvic RT in patients with endometrial carcinoma. In 2 of these trials, the patients were not formally staged (Postoperative Radiation Therapy in Endometrial Carcinoma [PORTEC-1], Aalders).<sup>191,192</sup> In the third trial (ASTEC/EN.5), only 50% of the patients were thoroughly staged as part of a companion surgical protocol.<sup>66,193</sup> However, formal surgical staging was mandated for all patients in the fourth trial (GOG 99).<sup>194</sup> Note that these trials used the older staging system (ie, before 2009).

The PORTEC-1 trial suggested that external-beam pelvic RT provides a therapeutic benefit in selected patients with uterine-confined disease.<sup>191,195</sup> Although RT significantly decreased locoregional recurrence, it did not increase OS.<sup>196</sup> The Aalders' randomized trial found that RT reduced vaginal (ie, locoregional) recurrences but did not reduce distant metastases or improve survival.<sup>192</sup> A recent pooled randomized trial (ASTEC/EN.5) suggested that adjuvant pelvic RT alone did not improve either relapse-free survival (ie, PFS) or OS in patients with intermediate-risk or high-risk early-stage endometrial cancer, but there

was a small improvement in pelvic control.<sup>193</sup> However, the ASTEC/EN.5 study is very controversial; 51% of the patients in the ASTEC observation group received vaginal brachytherapy.<sup>70,197</sup> The Keys' trial (GOG 99) showed that adjuvant pelvic RT improved locoregional control and relapse-free interval (ie, PFS), without OS benefit.<sup>194</sup> Both the GOG 99 and PORTEC-1 trials revealed that most of the initial recurrences for patients with initial uterine-confined tumors were limited to the vagina, prompting the increasing use of vaginal brachytherapy alone as adjunctive treatment.<sup>194,198,199</sup>

To help select a patient population who may benefit from adjuvant RT, the GOG 99 and PORTEC trials defined risk factors for women at high-intermediate risk (HIR) for recurrence.<sup>191,194</sup> These risk factors include age, in addition to deep myometrial invasion, grade, and LVSI. In GOG 99, women younger than 50 years had to have all 3 histologic risk factors to be considered HIR.<sup>194</sup> If they were 50 to 70 years, they were considered HIR if they had 2 histologic risk factors. Women 70 years or older were defined as HIR if they also had one risk factor. In PORTEC-1, women had to have 2 of 3 risk factors (ie, age >60 years, deep myometrial invasion, grade 3 histology) to be considered at HIR for recurrence.<sup>191,198</sup>

Due to concerns about potential toxicity of external-beam pelvic RT, the role of vaginal brachytherapy alone in uterine-confined disease has been evaluated. PORTEC-2 randomly assigned patients to external-beam pelvic RT versus vaginal brachytherapy alone in uterine-confined disease. PORTEC-2 showed excellent and equivalent vaginal and pelvic control rates with both adjuvant radiation approaches and no difference in OS.<sup>200</sup> Given that vaginal brachytherapy is associated with significantly less toxicity than pelvic RT, vaginal brachytherapy alone is a reasonable choice for most patients with uterine-confined endometrial cancer who are deemed candidates for adjuvant radiotherapy.<sup>198-207</sup> The use of vaginal brachytherapy and/or whole pelvic RT should be carefully tailored to a

patient's pathologic findings. Both PORTEC-1 and PORTEC-2 specifically excluded patients with 1998 FIGO stage 1C and grade 3 endometrial carcinoma (2009 FIGO stage IB, grade 3);<sup>54</sup> thus, the use of adjuvant brachytherapy alone in the highest risk subset remains undetermined.

A recent trial (GOG 249) examined vaginal cuff brachytherapy and carboplatin/paclitaxel therapy (brachy+chemo) versus pelvic EBRT only in patients with high-risk, uterine-confined endometrial carcinoma (n= 601). Unlike PORTEC-2, GOG 249 reported significantly increased rates of nodal recurrence (primarily pelvic) in the brachy+chemo arm versus the pelvic EBRT arm. No significant between-group differences in vaginal or distant recurrence rates were observed. However there were more extrvaginal pelvic failures in the brachy+chemo arm. At a median follow-up of 53 months, 3-year recurrence-free survival was 82% for both treatment arms; 3-year OS was 88% for the brachy+chemo cohort and 91% for the pelvic EBRT cohort. Acute toxicity was more common and severe for patients receiving brachytherapy with chemotherapy. No differences in late-onset toxicities were observed.<sup>208</sup>

Analysis of pooled data from PORTEC-1 and PORTEC-2 ranked the predictive power of multiple variables on patient outcomes examined in these trials. Patient age, tumor grade, and LVSI were highly predictive for locoregional relapse (LRR), distant relapse (DR), OS, and DFS, and treatment given (EBRT versus vaginal brachytherapy) was predictive for LRR and DFS.<sup>181</sup> The benefit of adjuvant EBRT in the highest risk spectrum of uterine-confined disease remains controversial. Most NCCN Panel Members feel that patients with deeply invasive grade 3 tumors should receive adjuvant treatment. Two large retrospective SEER analyses of women with endometrial cancer found that adjuvant RT improved OS in those with high-risk disease.<sup>209,210</sup> In a meta-analysis of randomized trials, a subset analysis found that adjuvant pelvic RT for stage I disease was associated with a trend towards a survival advantage

in the highest-risk spectrum (eg, those with 1988 FIGO stage IC grade 3) but not in lower risk patients; however, other reviews have shown conflicting results.<sup>202,211-215</sup>

Recently, results were published from a long-term follow-up study (median 20.5 years) of 568 patients with early-stage endometrial carcinoma who were enrolled in the Aalders trial. The study compared long-term outcomes in women who received vaginal brachytherapy plus EBRT versus vaginal brachytherapy alone. The findings suggested no statistical difference in OS between the study groups, and in this cohort, patients younger than 60 years of age who received EBRT had increased incidence of secondary cancers and subsequent higher mortality rates.<sup>202</sup>

### *Adjuvant Systemic Therapy*

Patients with deeply invasive, grade 3, uterine-confined disease (2009 FIGO stage IB, grade 3 [formerly 1988 FIGO stage IC, grade 3]) have a relatively poor prognosis. Despite adjuvant therapy with pelvic RT, a significant number of patients continue to have an appreciable risk of distant metastases.<sup>194,195</sup> Therefore, some clinicians suggested that adding systemic therapy to adjuvant RT may provide added therapeutic benefit (ie, decrease in distant metastases).<sup>183,216</sup> Studies have evaluated the role of systemic therapy in highest risk uterine-confined disease.<sup>216,217</sup> PFS is improved with adjuvant sequential chemotherapy/RT.<sup>216</sup> However, the NCCN panel feels that adjuvant systemic therapy is a category 2B recommendation in this setting because an OS advantage has not been shown.<sup>216</sup> We await final results from GOG 249.

Carboplatin/paclitaxel is the preferred regimen in the adjuvant setting for high-risk uterine confined disease.<sup>218-220</sup>

### *Advanced Stage/Extrauterine Disease*

There is a consensus that patients with documented extrauterine disease are at increased risk for recurrence and need adjuvant therapy; however,

the optimal form of adjuvant therapy has yet to be determined.<sup>221-223</sup> Patients with extrauterine disease confined to the lymph nodes or the adnexa may be treated with pelvic or extended-field RT alone.<sup>224</sup> However, systemic therapy is regarded as the foundation of adjuvant therapy for patients with extrauterine disease.

For stage IIIA to IIIC disease, the recommended treatment options are systemic therapy and/or EBRT with (or without) vaginal brachytherapy. For stage IVA/IVB disease systemic therapy forms the mainstay of treatment and can be combined with EBRT and/or vaginal brachytherapy.

Previously, whole abdominal RT was used for carefully selected patients deemed at risk for peritoneal failure, and RT appeared to have provided therapeutic benefit in retrospective studies.<sup>225,226</sup> A randomized phase III GOG (122) trial assessed optimal adjuvant therapy for patients with endometrial cancer who had extrauterine disease. In this trial, patients with stage III and intra-abdominal stage IV disease who had minimal residual disease were randomly assigned to whole abdominopelvic RT versus 7 cycles of combined doxorubicin (60 mg/m<sup>2</sup>) and cisplatin (50 mg/m<sup>2</sup>) treatment, with an additional cycle of cisplatin (AP). This GOG trial reported that AP chemotherapy improved PFS and OS when compared with whole abdominopelvic RT; however, acute toxicity (eg, peripheral neuropathy) was greater in the AP chemotherapy arm.<sup>178</sup>

The GOG 122 study established the role of adjuvant multiagent systemic chemotherapy for curative intent in patients with extrauterine disease. Thus, in the NCCN Guidelines, systemic therapy forms the established framework of adjuvant therapy for patients with stage III or IV disease. Whole abdominal RT as a single modality (as used in GOG 122) is considered inferior to chemotherapy and is no longer recommended. For the purposes of these guidelines, whole abdominal radiotherapy is not considered to be tumor-directed RT (see *Principles of Radiation Therapy* in the NCCN Guidelines for Uterine Neoplasms). Multimodality therapy is

now the basis of randomized trials evaluating therapy (eg, GOG 258, PORTEC-3).

Recurrences were frequent in both treatment arms of GOG 122, occurring in the pelvis and abdomen. Approximately 52% of patients with advanced endometrial carcinoma had recurrences, indicating the need for further therapeutic improvement in this high-risk patient population.<sup>178</sup> A study found that combined modality adjuvant therapy (using chemotherapy and tumor-directed RT) may provide a therapeutic benefit when compared with other sequencing modalities (chemotherapy followed by RT or vice versa).<sup>180,227,228</sup>

A follow-up study evaluated the role of chemotherapy “intensification” for this patient population. The GOG 184 trial assessed combination chemotherapy (cisplatin and doxorubicin with [or without] paclitaxel) with more limited radiation fields (involved-field radiation either to the pelvis or to the pelvis plus para-aortic nodes). Results indicate that the 3-drug regimen did not improve survival when compared with the 2-drug regimen after 3 years of follow-up and that the more intensive chemotherapy resulted in greater toxicity (eg, hematologic toxicity, sensory neuropathy, myalgia).<sup>179</sup>

Adjuvant therapy options were compared in a multicenter retrospective analysis of 265 patients with optimally resected stage IIIC endometrial carcinoma. Compared with patients receiving adjuvant RT or adjuvant RT plus chemotherapy, patients who received adjuvant chemotherapy had a 2.2 fold increased risk of recurrence and a 4.0 fold increased risk of death.<sup>223</sup> In a retrospective review of 116 patients with stage IIIC endometrial cancer, adjuvant RT significantly improved OS in patients with endometrioid histology, high-grade tumors, and positive para-aortic lymph nodes. Conversely, patients with low-grade tumors and non-endometrioid histology that received RT had similar OS compared with those who did not.<sup>229</sup> In a multicenter retrospective review of 73 patients with stage IIIA

endometrial carcinoma, surgery followed by both chemotherapy and radiation therapy provided the highest 5-year OS.<sup>230</sup> A prospective study of 122 patients with fully resected locally advanced disease suggested a potential benefit of adjuvant chemoradiation followed by chemotherapy, with an estimated 5-year PFS and OS of 73% and 84%.<sup>231</sup> The role of adjuvant RT with systemic therapy for treating high-risk endometrial carcinoma remains an area of active investigation (eg, GOG 258, PORTEC-3).

### Radiotherapy Principles

RT has been a widely used modality in the treatment of patients with endometrial cancer; it clearly improves locoregional control.

*Tumor-directed RT* refers to RT directed at sites of known or suspected tumor involvement and may include EBRT and/or vaginal brachytherapy.<sup>184</sup> RT is described in detail in the algorithm, including target areas and doses for pelvic RT and brachytherapy (see *Principles of Radiation Therapy* in the NCCN Guidelines for Uterine Neoplasms).

Although adjuvant RT is typically not associated with high rates of severe morbidity,<sup>232</sup> studies have focused on subtle effects on quality of life (eg, diarrhea, bowel symptoms) that deserve further investigation.<sup>203,205,233</sup> In the PORTEC-2 trial, vaginal brachytherapy was associated with better quality of life when compared with EBRT without a significant detriment to outcome.<sup>203</sup> Therefore, many patients who were previously treated with adjuvant EBRT are now appropriately treated with vaginal brachytherapy; this recommendation is reflected in the NCCN Guidelines. Patients treated with RT are prone to vaginal stenosis, which can impair sexual function. Women can use vaginal dilators to prevent or treat vaginal stenosis. Dilator use can start 2 to 4 weeks after RT is completed and can be used indefinitely (<http://www.mskcc.org/cancer-care/patient-education/resources/improving-your-vaginal-health-after-radiation-therapy>).

### Post-Treatment Surveillance

The recommended post-treatment surveillance protocol for endometrial cancer is shown in the algorithm (see *Surveillance* in the NCCN Guidelines for Endometrial Carcinoma).<sup>35</sup> These recommendations recognize that the value of intensive surveillance has not been demonstrated in this disease; therefore, ancillary testing is not recommended.<sup>234,235</sup>

Patients with clinical stage I and stage II endometrial cancer have a recurrence rate of approximately 15%;<sup>235-238</sup> 50% to 70% of these patients are symptomatic. For most patients, disease recurs within 3 years of initial treatment. Because most recurrences are symptomatic, all patients should receive verbal and written information regarding the symptoms of recurrent disease.<sup>235</sup> Patients with bleeding (vaginal, bladder, or rectal), decreased appetite, weight loss, pain (in the pelvis, abdomen, hip, or back), cough, shortness of breath, and swelling (in the abdomen or legs) should seek prompt evaluation and not delay until the next scheduled appointment.

As clinically indicated, imaging may be helpful in the detection of recurrence. The panel recommends imaging based on symptomatology and clinical concern for metastatic disease. In patients with treated stage III/IV disease, chest/abdominal/pelvic CT is an optional recommendation every 6 months during the first 3 years of surveillance, and every 6 to 12 months for an 2 additional years.

For suspected recurrence or metastasis, several imaging recommendations were added in 2017. Abdominal/pelvic CT and/or chest CT is recommended based on symptoms or physical exam findings suspicious for recurrence or metastasis. Additional imaging considerations include whole body PET/CT in select patients who may be candidates for surgery/locoregional therapy and/or pelvic MRI for patients who retain their uterus.

In the absence of recurrence, post-treatment surveillance provides psychosocial reassurance and improves quality of life for patients and their families. Health maintenance has been incorporated into the follow-up schedule (eg, blood pressure determination, breast examination, mammography as clinically indicated, stool guaiac test, immunizations). Patients should receive counseling and education regarding lifestyle, obesity, exercise, smoking cessation, sexual health, nutrition, and potential late or long-term effects of treatment (see the NCCN Guidelines for Survivorship, the NCCN Guidelines for Smoking Cessation, and <http://www.cancer.org/treatment/survivorshipduringandaftertreatment/index>).<sup>233,239-241</sup> Other health problems that often coexist in patients with endometrial cancer can also be evaluated during follow-up.

Given the lack of prospective studies regarding the optimal frequency of post-treatment follow-up, the NCCN Panel believes that the algorithm represents a reasonable surveillance scheme. The use of vaginal cytology is no longer recommended for asymptomatic patients consistent with the SGO guidelines.<sup>234,235,238,242</sup> Patients with stage I endometrial cancer have a low risk of asymptomatic vaginal recurrence (2.6%), especially after adjuvant brachytherapy, and vaginal cytology is not independently useful for detecting recurrences in this group of patients.<sup>234,243</sup> A recent multi-institutional review examined the utility of various surveillance methods in 254 patients with high-grade disease, revealing that symptoms led to the detection of the most recurrences (56%), followed by physical exam (18%), surveillance CT (15%), CA-125 (10%), and vaginal cytology (1%).<sup>244</sup>

### Hormone Replacement Therapy for Hypoestrogenism

After BSO, hypoestrogenism is associated with hot flashes, mood lability, vaginal dryness, pelvic soft tissue atrophy, osteoporosis, and an increased risk of cardiovascular disease. In postmenopausal women, estrogen replacement therapy was believed to reduce or reverse some of these

signs and symptoms. However, women who have had BSO for endometrial adenocarcinoma have usually been denied estrogen replacement therapy for fear of inducing a higher relapse rate, because this cancer has historically been considered an estrogen-linked malignancy.<sup>245,246</sup> As such, estrogen replacement therapy for such patients remains controversial.

However, it has never been proven that relapse rates are higher in endometrial cancer patients who receive estrogen replacement therapy after hysterectomy. Several retrospective trials of estrogen replacement after treatment of early-stage endometrial cancer have shown no increase in tumor recurrence or cancer-related deaths.<sup>247-249</sup> In women with stage I to II endometrial cancer who had hysterectomy, a randomized trial of estrogen replacement therapy versus placebo did not find an increased rate of recurrence or new malignancy; the median follow-up was 35.7 months.<sup>250</sup> However, estrogen replacement trials in postmenopausal females without a history of malignancy have demonstrated a significantly increased risk for breast cancer.<sup>251</sup>

Initially, the Women's Health Initiative (WHI) Estrogen-Alone Trial in women who had hysterectomy (n = 10,739) reported that the risk of breast cancer and cardiovascular disease (eg, stroke) were increased and that estrogen replacement therapy was of concern; thus, the trial was stopped.<sup>252</sup> However, recent long-term follow-up data from this trial suggest that the risk from estrogen-alone replacement therapy (without progesterone) may not be as high in younger women (<60 years) who have had hysterectomy.<sup>253</sup>

The NCCN panel agrees that estrogen replacement therapy is a reasonable option for patients who are at low risk for tumor recurrence, but initiating such therapy should be individualized and discussed in detail with the patient.<sup>254,255</sup> If adjuvant treatment is carried out, there should be a 6- to 12-month waiting period before initiation of hormone replacement therapy,

and participation in clinical trials is strongly encouraged. Selective estrogen-receptor modulators (SERMs) may prove to be attractive options for hormone replacement therapy.<sup>256,257</sup> Long-term comparisons between conjugated estrogens and SERMs for hormone replacement therapy are needed. Non-hormonal therapy may be considered in patients who are deemed poor candidates for hormone replacement therapy (eg, smokers, history of breast cancer, history of multiple strokes).<sup>258,259</sup>

### Treatment of Recurrent or Metastatic Disease

#### Localized Disease

Patients with local or regional recurrences can be evaluated for further treatment (see *Clinical Presentation* in the NCCN Guidelines for Endometrial Carcinoma). For recurrences confined to the vagina or the pelvis alone, second-line treatment (typically with RT and/or surgery or systemic therapy) can be effective. For patients with no prior RT exposure at the recurrence site or previous brachytherapy only, the panel recommends RT plus brachytherapy, or surgery. Isolated vaginal recurrences treated with RT have good local control and 5-year survival rates of 50% to 70%.<sup>260-262</sup> Prognosis is worse if there is extravaginal extension or pelvic lymph node involvement.<sup>261</sup> After RT, it is unusual for patients to have recurrences confined to the pelvis. The management of such patients remains controversial. For patients previously treated with brachytherapy only at the recurrence site, surgery with (or without) intraoperative RT (IORT) is recommended (category 3 for IORT). For patients previously treated with EBRT at the recurrence site, recommended therapy for isolated relapse includes: 1) surgery with (or without) IORT (category 3 for IORT); and/or 2) systemic therapy with (or without) palliative RT. In selected patients, radical surgery (ie, pelvic exenteration) has been performed with reported 5-year survival rates approximating 20%.<sup>263-266</sup>

Treatment for para-aortic or common iliac lymph node invasion and for upper abdominal or peritoneal recurrences is shown in the algorithm (see *Additional Therapy* in the NCCN Guidelines for Endometrial Carcinoma). However, for gross upper abdominal residual disease, more aggressive treatment for relapse is recommended, as outlined for disseminated metastases in *Therapy for Relapse* in the NCCN Guidelines for Endometrial Carcinoma. For resectable isolated metastases, consider surgical resection and/or RT, or ablative therapy. Providers can also consider systemic therapy (category 2B). Further recurrences or disease not amenable to local therapy are treated as disseminated metastases. Palliative care measures should also be considered in management of patients with systemic disease (see the NCCN Guidelines for Palliative Care and <http://emedicine.medscape.com/article/270646-overview>).

#### Systemic Disease

For patients with low-grade, asymptomatic, and hormone receptor–positive disseminated metastases, options include hormone therapy followed by systemic therapy on progression. Symptomatic, higher grade, or large-volume metastases can be treated with systemic therapy with (or without) palliative RT. For persistent progression of disseminated metastases, best supportive care or enrollment in a clinical trial is recommended.

#### Hormonal Therapy

The role of hormonal therapy in recurrent or metastatic cancer has been primarily evaluated in patients with endometrioid histologies only. Hormonal therapy is also used for selected patients with ESS (see section on *Uterine Sarcomas* in this Discussion). Hormonal agents for treating metastatic disease include megestrol with alternating tamoxifen, progestational agents alone, aromatase inhibitors, tamoxifen alone, or fulvestrant.<sup>165-167,267-269</sup> No particular drug, dose, or schedule has been found to be superior. The main predictors of response in the treatment of

metastatic disease are well-differentiated tumors, expression of ER/PR receptors, a long disease-free interval, and the location and extent of extrapelvic (particularly pulmonary) metastases.

For asymptomatic or low-grade disseminated metastases, hormonal therapy with progestational agents has shown good responses, particularly in patients with ER/PR-positive disease.<sup>169,270-272</sup> Tamoxifen has a 20% response rate in disease that does not respond to standard progesterone therapy.<sup>273,274</sup> Tamoxifen has also been combined with progestational agents; however, a few patients had grade 4 thromboembolic events with this combination regimen.<sup>165,267,275</sup> In some patients, aromatase inhibitors (eg, anastrozole, letrozole) may be substituted for progestational agents or tamoxifen.<sup>168,169,272,276</sup>

Other hormonal modalities have not been well-studied, and adjuvant therapy with hormonal agents has not been compared with cytotoxic agents.<sup>169,277</sup> If disease progression is observed after hormonal therapy, cytotoxic chemotherapy can be considered. However, clinical trials or best supportive care (see the NCCN Guidelines for Palliative Care) are appropriate for patients with disseminated metastatic recurrence who have a poor response to hormonal therapy and chemotherapy.

### Systemic Therapy

Chemotherapy for endometrial cancer has been extensively studied.<sup>278,279</sup> Based on the current data, multiagent chemotherapy regimens are preferred for metastatic, recurrent, or high-risk disease, if tolerated. Single-agent therapy can also be used (see *Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease* in the NCCN Guidelines for Endometrial Carcinoma). Recommended multiagent chemotherapy regimens include carboplatin/paclitaxel, cisplatin/doxorubicin, cisplatin/doxorubicin/paclitaxel, carboplatin/docetaxel, carboplatin/paclitaxel/bevacizumab, ifosfamide/paclitaxel (for

carcinosarcoma, category 1), cisplatin/ifosfamide (for carcinosarcoma), carboplatin/paclitaxel/trastuzumab (for HER2-positive serous carcinoma), and everolimus/letrozole (for endometrioid histologies). See *Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease* in the NCCN Guidelines for Endometrial Carcinoma.

A phase III randomized trial (GOG 177) compared 2 combination chemotherapy regimens in women with advanced/metastatic or recurrent endometrial carcinoma. The 273 women were randomly assigned to 1) cisplatin, doxorubicin, and paclitaxel; or 2) cisplatin and doxorubicin. The 3-drug regimen was associated with improved survival (15 vs. 12 months,  $P < .04$ ) but with significantly increased toxicity (ie, peripheral neuropathy); therefore, it is not widely used.<sup>280,281</sup> These regimens are now category 2A in the NCCN Guidelines, because most panel members feel that carboplatin/paclitaxel is a less toxic regimen (see *Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease* in the NCCN Guidelines for Endometrial Carcinoma). The response rates with other multiagent chemotherapy have ranged from 31% to 81% but with relatively short durations. The median survival for patients in such trials remains approximately 1 year.<sup>278,279</sup>

Carboplatin and paclitaxel is an increasingly used regimen for advanced/metastatic or recurrent endometrial cancer; the response rate is about 40% to 62%, and OS is about 13 to 29 months.<sup>282-285</sup> A phase III trial (GOG 209) compared carboplatin and paclitaxel versus cisplatin, doxorubicin, paclitaxel, and filgrastim (granulocyte-colony stimulating factor).<sup>282</sup> Trial data presented at a national meeting show that oncologic outcomes are similar, but the toxicity and tolerability profile favor carboplatin/paclitaxel. Thus, the carboplatin/paclitaxel regimen is now the preferred approach for many patients. For patients in whom paclitaxel is contraindicated, docetaxel can be considered in combination with carboplatin.<sup>286</sup>

A phase II trial initially examined the addition of bevacizumab to carboplatin and paclitaxel among 15 patients with advanced or recurrent endometrial carcinoma.<sup>287</sup> Although this study was closed early due to the initiation of a national trial, a retrospective analysis was performed to include data from an additional 27 patients who had received carboplatin/paclitaxel/bevacizumab for advanced or recurrent disease.<sup>288</sup> Collective median PFS was 20 months with a median OS of 56 months. An overall response rate of 82.8% was noted, with an 87.5% response rate among the subset of 8 patients who received this triplet as second-line therapy after carboplatin/paclitaxel.<sup>288</sup>

Everolimus combined with letrozole is under investigation for recurrent disease with positive preliminary findings in disease of endometrioid histology. In the preliminary results, the clinical benefit rate and objective response rate among 35 evaluable patients was 40% and 32%, respectively.<sup>289</sup>

If multiagent chemotherapy regimens are contraindicated, then single-agent therapy options include paclitaxel, albumin-bound paclitaxel, cisplatin, carboplatin, doxorubicin, liposomal doxorubicin, topotecan, and docetaxel (category 2B for docetaxel) (see *Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease* in the NCCN Guidelines for Endometrial Carcinoma).<sup>169,290-292</sup> When single agents are used as first-line treatment, responses range from 21% to 36%.<sup>272,293</sup> When single agents are used as second-line treatment, responses range from 4% to 27%; paclitaxel is the most active in this setting.<sup>293</sup> Some oncologists have used liposomal doxorubicin, because it is less toxic than doxorubicin; the response rate of liposomal doxorubicin is 9.5%.<sup>294</sup> Docetaxel is recommended for use as a single agent; however, it is a category 2B recommendation because some panel members would not use docetaxel because it is less active (7.7% response rate) than other agents.<sup>164,295</sup>

In the advanced endometrial cancer cohort (n = 24) of the phase Ib KEYNOTE-028 trial, durable antitumor responses were noted in a small subset of patients with programmed death ligand 1 (PD-L1) positive tumors (3 partial response, 3 stable disease).<sup>296</sup> Studies have also indicated that MMR-deficient (dMMR) tumors are sensitive to programmed death receptor-1 (PD-1) blockade.<sup>297-299</sup> Results were recently published from a study of patients with dMMR tumors of various disease sites. Among patients with dMMR endometrial carcinoma who received pembrolizumab (n = 15), the objective response rate was 52% and the disease control rate was 73% (3 complete response, 5 partial response, and 3 stable disease).<sup>297</sup> The FDA expanded pembrolizumab approval in 2017 to include treatment of unresectable or metastatic, MSI-high (MSI-H), or dMMR solid tumors that have progressed following prior treatment and that have no satisfactory alternative treatment options.<sup>299</sup> The panel voted to include pembrolizumab as a treatment option for MSI-H/dMMR endometrial tumors and recommends that recurrent endometrial tumors be tested for MSI-H or dMMR if not done previously.

New biologic and molecular therapies for the treatment of recurrent or metastatic endometrial carcinoma are being assessed in clinical trials.<sup>164,300</sup> Bevacizumab was shown to have a 13.5% response rate and OS rate of 10.5 months in a phase II trial for persistent or recurrent endometrial cancer.<sup>301</sup> Temezirolimus has been used as first-line or second-line therapy for recurrent or metastatic endometrial cancer and has a partial response rate of 4% in second-line therapy.<sup>302</sup> Based on these studies, the NCCN panel considers bevacizumab or temsirolimus as appropriate single-agent biologic therapy for patients who have progressed on previous cytotoxic chemotherapy.<sup>300-303</sup>

### Uterine Serous Carcinomas, Clear Cell Carcinomas, and Carcinosarcomas

#### Overview

Uterine serous carcinomas, clear cell carcinomas, carcinosarcomas, and undifferentiated/dedifferentiated carcinomas are considered more aggressive histologic variants of malignant epithelial tumors, with a higher incidence of extrauterine disease at presentation.<sup>304-311</sup> Carcinosarcomas are aggressive tumors that are staged as high-grade endometrial cancer.<sup>312,313</sup> Serous carcinomas, clear cell carcinomas, carcinosarcomas, and undifferentiated/dedifferentiated carcinomas are all considered high-risk histologies and high grade by default, although they are staged using the same FIGO/AJCC staging system as endometrial cancers (see Table 1).<sup>56</sup>

Pathologists now believe that carcinosarcomas (also known as MMTTs) are metaplastic carcinomas and not uterine sarcomas; therefore, carcinosarcomas are included in the high-risk malignant epithelial tumors section of the NCCN Guidelines (see *Serous Carcinoma, Clear Cell Carcinoma, or Carcinosarcoma* in the NCCN Guidelines for Endometrial Carcinoma).<sup>308,311,314,315</sup> Even patients with apparent early-stage disease may have distant metastases. Thus, fertility-sparing therapy is not recommended for these aggressive tumors. If done, SLN mapping should proceed with particular caution.

Patients with uterine serous carcinoma, clear cell carcinoma, carcinosarcoma, or undifferentiated/dedifferentiated carcinomas may present with pelvic masses, abnormal cervical cytology, or ascites in addition to postmenopausal bleeding. Both the NCCN panel and the SGO recommend that CA-125 and MRI or chest/abdominal/pelvic CT may be useful before surgery to assess if extrauterine disease is present; PET may also be useful.<sup>304</sup> Patterns of failure often mimic those of ovarian cancer.

#### Treatment

Multimodality therapy is typically recommended for these histologically aggressive tumors. Primary treatment includes TH/BSO with surgical staging, peritoneal lavage for cytology, omental and peritoneal biopsies, and consideration of maximal tumor debulking for gross disease (see *Principles of Evaluation and Surgical Staging* in the NCCN Guidelines for Endometrial Carcinoma).<sup>316</sup> Minimally invasive surgery is the preferred approach when technically feasible.<sup>317-321</sup>

Adjuvant therapy is highly individualized.<sup>322-329</sup> For patients with stage IA without myometrial invasion, options include: 1) chemotherapy with (or without) vaginal brachytherapy (preferred approach); 2) observation if no residual serous or clear cell carcinoma in hysterectomy specimen; or 3) EBRT with (or without) vaginal brachytherapy.<sup>330,331</sup> For all other patients with more advanced disease, systemic therapy with (or without) tumor-directed RT is the preferred option.<sup>306,323,327,332</sup> Adjuvant platinum/taxane-based therapy appears to improve survival in patients with uterine serous carcinoma and clear cell carcinoma, whereas ifosfamide/paclitaxel (category 1) is recommended for carcinosarcomas (see *Uterine Serous Carcinomas, Clear Cell Carcinomas, and Carcinosarcomas* in this Discussion and *Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease* in the NCCN Guidelines for Endometrial Carcinoma).<sup>304-306,333-335</sup>

Whole abdominopelvic RT with (or without) vaginal brachytherapy is no longer recommended as a primary treatment option for patients with advanced disease, because the NCCN panel no longer feels that routine use of whole abdominal RT is appropriate.<sup>178,332,336</sup> Chemotherapy with (or without) RT appears to be more effective than RT alone.<sup>323</sup> Data are conflicting regarding the rate of abdominal recurrence in these patients.<sup>332,337-341</sup> Whole abdominal radiotherapy is not considered to be tumor-directed RT (see *Principles of Radiation Therapy* in the NCCN

Guidelines for Uterine Neoplasms). As previously mentioned, *tumor-directed RT* refers to RT directed at sites of known or suspected tumor involvement and may include EBRT with (or without) vaginal brachytherapy. In general, tumor-directed EBRT is directed to the pelvis with (or without) the para-aortic region.

Several studies have examined treatment paradigms for uterine serous or clear cell carcinoma. A recent phase II trial in patients with papillary serous carcinoma suggested favorable outcomes with concurrent pelvic RT and paclitaxel followed by additional rounds of adjuvant paclitaxel,<sup>342</sup> indicating the potential benefits of combined modality therapy. Retrospective data were reviewed from 279 patients with serous or clear cell carcinoma who were treated at high-volume cancer centers. Adjuvant treatment (RT, systemic therapy, or chemoradiation) was associated with improved OS in stages IB-II disease but not stage IA disease (HR, 0.14; 95% CI, 0.02–0.78;  $P = .026$ ).<sup>317</sup> Additionally, survival outcomes did not differ based upon surgical approach (robotic laparoscopy versus laparotomy).

Two multi-institutional retrospective reviews examined the impact of adjuvant therapy (vaginal brachytherapy,  $n = 103$ ; adjuvant pelvic radiation or chemotherapy,  $n = 115$ ) in patients with stage 1A uterine papillary serous carcinoma. In both cohorts, patients undergoing surgical staging/lymphadenectomy had greater PFS and OS than unstaged patients.<sup>343,344</sup> Vaginal brachytherapy reduced the vaginal recurrence rate but did not impact PFS or OS.<sup>343</sup> In unstaged patients, chemotherapy or pelvic RT were associated with greater PFS and OS, but no survival benefits were observed for adjuvant treatment in surgically staged patients.<sup>344</sup>

A recent randomized phase II study examined the addition of trastuzumab to carboplatin/paclitaxel for patients with advanced or recurrent human epidermal growth factor receptor 2 (HER2)/neu–positive uterine serous carcinoma.<sup>345</sup> Among patients with stage III/IV disease undergoing primary

treatment ( $n = 41$ ), median PFS was 17.9 months versus 9.3 months for the experimental and control arms, respectively ( $P = 0.013$ ). PFS for patients with recurrent disease ( $n = 17$ ) was 9.2 months versus 6.0 months ( $P = 0.003$ ). The addition of trastuzumab appeared to improve PFS without increasing overall toxicity.

For treating carcinosarcoma, ifosfamide was historically considered the most active single agent.<sup>334,346,347</sup> A phase III trial for advanced carcinosarcoma showed that the combination of ifosfamide and paclitaxel increased survival and was less toxic than the previously used cisplatin/ifosfamide regimen.<sup>334,348</sup> OS was 13.5 months with ifosfamide/paclitaxel versus 8.4 months with ifosfamide alone. Therefore, ifosfamide/paclitaxel is a category 1 recommendation in the NCCN Guidelines (see *Systemic Therapy for Recurrent, Metastatic, or High-Risk Disease* in the NCCN Guidelines for Endometrial Carcinoma).<sup>334,336</sup> However, the toxicity of ifosfamide has led to investigation of better-tolerated regimens. A phase II trial suggests that paclitaxel/carboplatin is also a useful regimen for carcinosarcoma (response rate, 54%).<sup>349</sup> A GOG trial is currently assessing ifosfamide/paclitaxel versus carboplatin/paclitaxel.<sup>312</sup> The panel now considers carboplatin/paclitaxel the preferred adjuvant therapy regimen for uterine-confined endometrial cancers, including carcinosarcoma.

Data regarding carcinosarcoma suggest that adjuvant pelvic radiotherapy decreases the rate of local recurrences when compared with surgery alone.<sup>350-355</sup> This local control improvement in some series correlates with an improvement in survival, although other data show that lymphadenectomy confers greater benefit.<sup>354-357</sup> A phase III randomized trial (GOG 150) in patients with carcinosarcoma of the uterus showed a trend towards a decreased mortality rate for patients receiving cisplatin/ifosfamide vs. whole-abdominal RT ( $P = .085$ ), although these did not reach statistical significance in this underpowered trial.<sup>336,341</sup> A recent

cohort study in women with early-stage MMT suggests that postoperative chemotherapy improves PFS compared to RT or observation.<sup>312</sup>

## Uterine Sarcomas

### Overview

In 2017, an estimated 4910 cases of uterine sarcomas are anticipated.<sup>358</sup> Uterine sarcomas are malignant mesenchymal tumors that include ESS, UUS, and uLMS (see *Uterine Sarcoma Classification* in the NCCN Guidelines for Uterine Sarcoma). According to a 2012 systematic review of data from 1970 to 2011, uLMS was the most common subtype (63%), followed by ESS (21%) and less common subtypes such as UUS.<sup>359</sup> Even rarer subtypes of malignant mesenchymal tumors that can occur in the uterus include adenosarcoma, rhabdomyosarcoma (RMS), and PEComa.<sup>360</sup> Carcinosarcomas were previously categorized and included in the sarcoma treatment algorithms until the mid-2000s, but are now considered and treated as high-grade epithelial tumors (carcinomas).<sup>308</sup> Screening for Lynch syndrome is not usually done for patients with malignant mesenchymal tumors.

ESSs are composed of cells resembling the endometrial stroma in the proliferative phase.<sup>360,361</sup> ESS displays a heterogenous mix of morphologic and genetic features. A significant proportion of these tumors (ie, up to half) harbor a *JAZF1-SUZ12* (formerly *JAZF1-JJAZ1*) gene fusion and present as lower grade, earlier stage tumors.<sup>362-365</sup> More recently, a higher-grade and more aggressively behaving ESS variant with a unique genetic rearrangement *YWHAE-FAM22A/B*, also known as *YWHAE-NUTM2A/B*, was identified.<sup>366,367</sup> These findings provided support for subdividing ESS into distinct low- and high-grade entities based on histopathology, clinical behavior, and patient outcomes. In light of new information, WHO released an updated (4<sup>th</sup>) edition of the *WHO Classification of Tumours of Female*

*Reproductive Organs*. The updated 2014 edition recognizes low-grade ESS and high-grade ESS as distinct histopathologic entities.<sup>368</sup>

Recent advances have expanded our understanding of the molecular features of these tumors, leading to the identification of genetic signatures that characterize some of the uterine sarcoma subtypes. At present, mesenchymal tumors are primarily diagnosed using histopathologic criteria, and the results of molecular studies are not used in routine pathologic evaluation. However, molecular analysis (eg, identification of characteristic translocations) can help classify difficult cases and provide future therapeutic targets.

### Staging and Treatment

When evaluating suspected uterine sarcomas, biopsy may be helpful but is less sensitive than for endometrial cancers. The diagnosis of ESS and uLMS is often made after hysterectomy. The previous FIGO/AJCC staging systems for endometrial cancer were not appropriate for staging ESS and uLMS; patients were often upstaged when using the older AJCC staging system.<sup>369</sup> A new staging system for ESS and uLMS from FIGO/AJCC took effect in 2009 accounting for the differences between uterine sarcomas and endometrial cancers.<sup>56,370</sup>

Confirmation of the type of mesenchymal malignancy by expert pathology review is critical. In addition, initial evaluation should include imaging of the chest/abdomen/pelvis by CT or combination MRI/CT. It is important to determine if the sarcoma is confined to the uterus or if extrauterine disease is present. Pelvic MRI can be used to evaluate local tumor extension or residual abnormality in cases where the uterus or adnexa were not resected or incompletely resected (ie, supracervical hysterectomy, myomectomy, possible tumor fragmentation, intraperitoneal morcellation). Whole-body PET/CT may be used to clarify ambiguous findings. If medically operable, then hysterectomy with (or without) BSO is

the initial treatment of choice for uterine sarcomas (see *Primary Treatment* in the NCCN Guidelines for Uterine Sarcoma).<sup>371</sup>

The panel recommends ER/PR testing to guide decisions regarding management of the ovaries, particularly in young premenopausal patients. In general, BSO is favored for low-grade ESS or tumors expressing ER/PR, although management of the ovaries may be individualized in reproductive-age patients.

Uterine sarcoma should be removed en bloc to optimize outcomes; intraperitoneal morcellation is contraindicated.<sup>142-145</sup> For incidental diagnoses of uterine sarcoma after hysterectomy, or in the case of a fragmented specimen, imaging is recommended and re-exploration can be considered. The ovaries may be preserved in selected patients with early-stage uLMS who wish to retain hormonal function.<sup>372</sup> Additional surgical resection should be individualized based on clinical scenarios and intraoperative findings. Lymphadenectomy is controversial.<sup>2,360,372-375</sup> High-grade uterine sarcomas tend to show hematogenous metastases to the lungs; lymph node metastases are uncommon.

For medically inoperable sarcomas, options include: pelvic EBRT with (or without) brachytherapy and/or systemic therapy.

### **Low-Grade Endometrial Stromal Sarcoma**

If there is no evidence of disease after primary surgery (TH/BSO) for stage I ESS, then observation can be considered (see *Additional Therapy* in the NCCN Guidelines for Uterine Sarcoma).<sup>373,374</sup> Recommended adjuvant therapy options for stage I ESS include observation (especially if menopause or prior BSO) or estrogen blockade (category 2B). Postoperative estrogen blockade is recommended for stages II to IV ESS. Adjuvant EBRT may be added for stage II-IVA (category 2B). Palliative RT may be added to estrogen blockade for patients with stage IVB disease.<sup>360,376,377</sup> Typical hormone therapy includes aromatase inhibitors

(preferred), megestrol acetate, or medroxyprogesterone acetate. Gonadotropin-releasing hormone [GnRH] analogs (category 2B) are also an option.<sup>360,372,378</sup> In 2014, tamoxifen was removed from the NCCN Guidelines for ESS because it is contraindicated in women diagnosed with ESS or ER/PR-positive uLMS.<sup>372,377-379</sup> Estrogen blockade is also recommended for ESSs that have recurred or are unresectable (see *Therapy for Relapse* in the NCCN Guidelines for Uterine Sarcoma).<sup>378</sup>

Case series of patients with ESS suggest long disease-free intervals in the absence of specific therapy and raise questions about the use of adjuvant RT.<sup>380</sup> Adjuvant radiotherapy in ESS has been demonstrated to reduce local recurrence rates but again with limited effect on survival.<sup>381,382</sup> Because of concerns about radiation exposure, frequent routine surveillance imaging is no longer recommended for asymptomatic young women after primary therapy for ESS.<sup>383</sup>

Although hormone therapy is recommended for low-grade ESS, studies have not yet determined the optimal therapeutic approach for high-grade ESS. However, due to the more aggressive nature of these tumors (eg, those with YWHA-E-FAM22 rearrangements), the NCCN panel has recommended that high-grade ESS be treated according to the algorithms in place for uLMS and UUS.

### **High-Grade Endometrial Stromal Sarcoma, Leiomyosarcoma, and Undifferentiated Uterine Sarcoma**

The role of adjuvant radiotherapy in nonmetastatic disease is controversial. Most available data are retrospective except for a phase III randomized trial.<sup>350</sup> Most retrospective studies of adjuvant RT suggest an improvement in local pelvic control but no appreciable or consistent improvement in OS, given the propensity of metastatic extrapelvic disease as a site of first or eventual recurrence.<sup>384-387</sup> In many series, the patients treated with adjuvant radiation presumably had higher-risk factors (eg, larger tumors, deeper myometrial invasion), thus biasing the data against

radiotherapy. However, a phase III randomized trial in stage I and II uterine sarcomas reported that postoperative pelvic radiotherapy did not improve OS for uLMS when compared with observation.<sup>350</sup> Therefore, routine postoperative RT is not recommended for stage I patients with uLMS and high-grade (undifferentiated) endometrial sarcoma.<sup>376</sup> If used in more advanced stages, adjuvant RT needs to be individualized and based on careful analysis of surgical pathologic findings.

The role of adjuvant systemic therapy is also poorly defined; however, adjuvant systemic therapy has been used because of the high risk of systemic relapse. Given the uncertainties regarding any adjuvant treatment for stage I uLMS and high-grade (undifferentiated) endometrial sarcoma, after complete resection options include: 1) observation; 2) systemic therapy (category 2B); or 3) estrogen blockade if ER positive may be considered. Because of the increased risk profile in patients with completely resected stage II and III uLMS and high-grade (undifferentiated) endometrial sarcoma, the panel believes that it is appropriate to consider adjuvant systemic therapy and/or EBRT (see *Additional Therapy* in the NCCN Guidelines for Uterine Sarcoma).<sup>388</sup> In patients with incompletely resected or metastatic disease, systemic therapy with (or without) palliative EBRT is generally recommended.

An ongoing phase III randomized trial (GOG 277) is assessing the role of postoperative adjuvant chemotherapy (ie, gemcitabine/docetaxel followed by doxorubicin) versus observation in patients with high-grade stage I and II uLMS.<sup>389</sup>

If systemic therapy is used for treating high-grade uterine sarcoma, preferred therapies include single-agent doxorubicin, and gemcitabine/docetaxel<sup>389-394</sup> combination therapy (see *Systemic Therapy* in the NCCN Guidelines for Uterine Sarcoma).<sup>360,361,395</sup> Doxorubicin is an active single agent for uLMS and is less toxic than combination regimens.<sup>360,395</sup>

Other recommended combination regimens include doxorubicin/ifosfamide, doxorubicin/dacarbazine, gemcitabine/dacarbazine, and gemcitabine/vinorelbine.<sup>347,391,396-398</sup> Other single-agent options (category 2A unless otherwise noted) can also be considered for advanced or metastatic disease, including dacarbazine, gemcitabine, epirubicin, ifosfamide, liposomal doxorubicin, pazopanib, temozolomide, trabectedin (for uLMS treated with a prior anthracycline-containing regimen), eribulin (category 2B), vinorelbine (category 2B), and docetaxel (category 3).<sup>360,361,390,396,397,399-415</sup> Aromatase inhibitors can be considered for ER/PR-expressing uLMS.<sup>416</sup> Dacarbazine was elevated to a category 2A recommendation (from a category 2B) in 2014 because this agent was used as the standard arm in several phase II trials.<sup>396</sup> In 2016, trabectedin and eribulin were both included in the guidelines.

Data indicate that trabectedin may be useful in patients who have exhausted standard chemotherapy.<sup>417-420</sup> Preliminary phase III data revealed a 2.7-month PFS benefit versus dacarbazine in metastatic liposarcoma or leiomyosarcoma that progressed after anthracycline-based therapy.<sup>421</sup> Follow-up subgroup analysis of patients with uLMS (n = 232) revealed PFS of 4.0 months for trabectedin versus 1.5 months for dacarbazine (HR, 0.57; 95% CI, 0.41–0.81; P = .0012).<sup>422</sup> However, OS did not differ significantly between the treatment arms (13.4 months for trabectedin vs. 12.9 months for dacarbazine; HR, 0.89; 95% CI, 0.65–1.24; P = .51). Following its October 2015 FDA approval, trabectedin was added to the guidelines as an option for unresectable or metastatic uLMS previously treated with an anthracycline-containing regimen.

Eribulin was included based on results from a phase III trial comparing the survival benefit of eribulin and dacarbazine in 452 patients with advanced leiomyosarcoma or adipocytic sarcoma.<sup>423</sup> Median OS was 13.5 and 11.5 months for eribulin and dacarbazine, respectively (HR, 0.77; 95% CI,

0.62–0.95;  $P = .017$ ). Eribulin was designated as category 2B upon panel review of the mature trial data.

### Post-Treatment Surveillance

The recommended post-treatment surveillance protocol for uterine sarcoma is depicted in the algorithm (see *Surveillance* in the NCCN Guidelines for Uterine Sarcoma). History and physical exam is recommended every 3 to 4 months for the first 2 to 3 years, and then every 6 to 12 months thereafter. Imaging surveillance should include chest/abdominal/pelvic CT every 3 to 6 months for the first 3 years and then every 6 to 12 months for the next 2 years. Depending on histology, grade, and initial stage, annual to biannual imaging can be considered for an additional 5 years. Abdominal/pelvic MRI and chest CT can also be considered, with PET/CT or other imaging as needed to clarify findings or upon clinical concern for metastasis.

Patients should receive education regarding the symptoms of recurrent disease. Patients with bleeding (vaginal, bladder, or rectal), decreased appetite, weight loss, pain (in the pelvis, abdomen, hip, or back), cough, shortness of breath, and swelling (in the abdomen or legs) should seek prompt evaluation and not delay until the next scheduled appointment. As clinically indicated, imaging may be helpful in the detection of recurrence. Patients should be educated regarding healthy lifestyle, obesity, exercise, smoking cessation, nutrition, and potential long-term and late effects of treatment (see the NCCN Guidelines for Survivorship, NCCN Guidelines for Smoking Cessation, and <http://www.cancer.org/treatment/survivorshipduringandaftertreatment/index>).<sup>239-241</sup> The panel also recommends patient education regarding sexual health, vaginal dilator use, and vaginal lubricants or moisturizers.

### Treatment of Recurrent or Metastatic Disease

The recurrence rate is high in uLMS (50%–70%).<sup>2</sup> The guidelines provide recommendations based on tumor resectability and patients' prior RT exposure (see *Therapy for Relapse* in the NCCN Guidelines for Uterine Sarcoma). Treatment recommendations are made according to the site and nature of the recurrence.

Local recurrences are classified as recurrence in the vagina/pelvis with imaging that is negative for distant metastatic disease. Surgical and RT treatment pathways are provided. The surgical pathway for treating local recurrence in patients without prior RT exposure includes the option of IORT (category 3 for IORT). Preoperative EBRT and/or systemic therapy are also options to consider. For residual disease following surgery in patients without preoperative RT, EBRT with (or without) brachytherapy and/or systemic therapy can be considered. Primary RT offers an alternative pathway for treating localized recurrence in patients without prior exposure. EBRT should be given along with the option of brachytherapy and systemic therapy. For both the surgical and RT treatment pathways, further adjuvant systemic therapy should be considered after initial treatment.

Patients with local recurrence who have had prior RT exposure can be treated with: 1) surgery with the option of IORT and/or systemic therapy (category 3 for IORT); 2) systemic therapy; or 3) selected reirradiation with EBRT and/or brachytherapy. A recent retrospective analysis of patients with ESS suggested that cytoreductive resection improved OS in patients with recurrent lesions.<sup>424</sup>

Systemic therapy with (or without) palliative EBRT or best supportive care is recommended for metastatic disease.<sup>395</sup> For patients with isolated metastases, surgical resection or other ablative therapy (eg, radiofrequency ablation, stereotactic body RT) may be appropriate.

Patients with uLMS who experience longer time to recurrence may have improved survival outcomes following metastasectomy.<sup>425</sup> Postoperative EBRT and/or systemic therapy can be considered. Systemic therapy and/or local therapy (tumor-directed EBRT or local ablative therapy) are reasonable options for patients with unresectable isolated metastases (see *Therapy for Relapse* in the NCCN Guidelines for Uterine Sarcoma).<sup>411,426-428</sup> For recurrent low-grade ESS, the first choice of systemic therapy is estrogen blockade.

### Drug Reactions

Virtually all drugs have the potential to cause adverse hypersensitivity reactions, either during or after the infusion.<sup>429</sup> In gynecologic oncology treatment, drugs that more commonly cause adverse reactions include carboplatin, cisplatin, docetaxel, liposomal doxorubicin, and paclitaxel. Most of these drug reactions are mild infusion reactions (ie, skin reactions, cardiovascular reactions, respiratory or throat tightness), but more severe allergic reactions (ie, life-threatening anaphylaxis) can occur.<sup>430-432</sup> In addition, patients can have mild allergic reactions or severe infusion reactions. Infusion reactions are more common with paclitaxel.<sup>433</sup> Allergic reactions (ie, true drug allergies) are more common with platinum agents (ie, carboplatin, cisplatin).<sup>433,434</sup>

Management of drug reactions is discussed in the NCCN Guidelines for Ovarian Cancer.<sup>433</sup> It is important to note that patients who have had severe life-threatening reactions should not receive the implicated agent again unless under the care of an allergist or expert in managing drug reactions. If a mild allergic reaction has previously occurred and it is appropriate to administer the drug again, a desensitization regimen should be used even if the symptoms have resolved; various desensitization regimens have been published and should be followed.<sup>435-437</sup> Patients must be desensitized with each infusion if they previously had a reaction.

Almost all patients can be desensitized (about 90%).<sup>429</sup> To maximize safety, it is prudent to desensitize patients in the intensive care unit.<sup>429</sup>

### References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin* 2018;68:7-30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29313949>.
2. D'Angelo E, Prat J. Uterine sarcomas: a review. *Gynecol Oncol* 2010;116:131-139. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19853898>.
3. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin* 2016;66:7-30. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26742998>.
4. Van den Bosch T, Coosemans A, Morina M, et al. Screening for uterine tumours. *Best Pract Res Clin Obstet Gynaecol* 2011. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22078749>.
5. Kitchener HC, Trimble EL. Endometrial cancer state of the science meeting. *Int J Gynecol Cancer* 2009;19:134-140. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19258955>.
6. Dinkelspiel HE, Wright JD, Lewin SN, Herzog TJ. Contemporary clinical management of endometrial cancer. *Obstet Gynecol Int* 2013;2013:583891. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23864861>.
7. Obermair A, Youlden DR, Young JP, et al. Risk of endometrial cancer for women diagnosed with HNPCC-related colorectal carcinoma. *Int J Cancer* 2010;127:2678-2684. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20533284>.
8. Katz VL. Diagnostic procedures. Imaging, endometrial sampling, endoscopy: indications and contraindications, complications. In: Katz VL, Lentz GM, Lobo RA, Gershenson DM, eds. *Comprehensive Gynecology*. 5th ed. Philadelphia, Pa: Mosby; 2007:chap 11.
9. Resnick KE, Hampel H, Fishel R, Cohn DE. Current and emerging trends in Lynch syndrome identification in women with endometrial cancer. *Gynecol Oncol* 2009;114:128-134. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19375789>.
10. Kwon JS, Scott JL, Gilks CB, et al. Testing women with endometrial cancer to detect Lynch syndrome. *J Clin Oncol* 2011;29:2247-2252. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21537049>.
11. Win AK, Lindor NM, Winship I, et al. Risks of colorectal and other cancers after endometrial cancer for women with Lynch syndrome. *J Natl Cancer Inst* 2013;105:274-279. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23385444>.
12. Lancaster JM, Powell CB, Kauff ND, et al. Society of Gynecologic Oncologists Education Committee statement on risk assessment for inherited gynecologic cancer predispositions. *Gynecol Oncol* 2007;107:159-162. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17950381>.
13. Meyer LA, Broaddus RR, Lu KH. Endometrial cancer and Lynch syndrome: clinical and pathologic considerations. *Cancer Control* 2009;16:14-22. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19078925>.
14. Bonnet D, Selves J, Toulas C, et al. Simplified identification of Lynch syndrome: a prospective, multicenter study. *Dig Liver Dis* 2012;44:515-522. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22480969>.
15. Walsh CS, Blum A, Walts A, et al. Lynch syndrome among gynecologic oncology patients meeting Bethesda guidelines for screening. *Gynecol Oncol* 2010;116:516-521. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20034658>.
16. Buchanan DD, Tan YY, Walsh MD, et al. Tumor mismatch repair immunohistochemistry and DNA MLH1 methylation testing of patients with endometrial cancer diagnosed at age younger than 60 years optimizes triage for population-level germline mismatch repair gene mutation testing. *J Clin Oncol* 2014;32:90-100. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24323032>.

17. Ferguson SE, Aronson M, Pollett A, et al. Performance characteristics of screening strategies for Lynch syndrome in unselected women with newly diagnosed endometrial cancer who have undergone universal germline mutation testing. *Cancer* 2014;120:3932-3939. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25081409>.
18. Goodfellow PJ, Billingsley CC, Lankes HA, et al. Combined Microsatellite Instability, MLH1 Methylation Analysis, and Immunohistochemistry for Lynch Syndrome Screening in Endometrial Cancers From GOG210: An NRG Oncology and Gynecologic Oncology Group Study. *J Clin Oncol* 2015;33:4301-4308. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26552419>.
19. Watkins JC, Yang EJ, Muto MG, et al. Universal Screening for Mismatch-Repair Deficiency in Endometrial Cancers to Identify Patients With Lynch Syndrome and Lynch-like Syndrome. *Int J Gynecol Pathol* 2017;36:115-127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27556954>.
20. Mills AM, Liou S, Ford JM, et al. Lynch syndrome screening should be considered for all patients with newly diagnosed endometrial cancer. *Am J Surg Pathol* 2014;38:1501-1509. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25229768>.
21. Smith RA, Cokkinides V, Brawley OW. Cancer screening in the United States, 2012: A review of current American Cancer Society guidelines and current issues in cancer screening. *CA Cancer J Clin* 2012. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22261986>.
22. Crispens MA. Endometrial and ovarian cancer in lynch syndrome. *Clin Colon Rectal Surg* 2012;25:97-102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23730224>.
23. Manchanda R, Saridogan E, Abdelraheim A, et al. Annual outpatient hysteroscopy and endometrial sampling (OHES) in HNPCC/Lynch syndrome (LS). *Arch Gynecol Obstet* 2012;286:1555-1562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22865035>.
24. Jarvinen HJ, Renkonen-Sinisalo L, Aktan-Collan K, et al. Ten years after mutation testing for Lynch syndrome: cancer incidence and outcome in mutation-positive and mutation-negative family members. *J Clin Oncol* 2009;27:4793-4797. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19720893>.
25. Schmeler KM, Lynch HT, Chen LM, et al. Prophylactic surgery to reduce the risk of gynecologic cancers in the Lynch syndrome. *N Engl J Med* 2006;354:261-269. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16421367>.
26. Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin* 2017;67:7-30. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28055103>.
27. Ueda SM, Kapp DS, Cheung MK, et al. Trends in demographic and clinical characteristics in women diagnosed with corpus cancer and their potential impact on the increasing number of deaths. *Am J Obstet Gynecol* 2008;198:218 e211-216. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18226630>.
28. Chan JK, Sherman AE, Kapp DS, et al. Influence of gynecologic oncologists on the survival of patients with endometrial cancer. *J Clin Oncol* 2011;29:832-838. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21263082>.
29. Benedetti Panici P, Basile S, Salerno MG, et al. Secondary analyses from a randomized clinical trial: age as the key prognostic factor in endometrial carcinoma. *Am J Obstet Gynecol* 2014;210:363 e361-363 e310. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24361787>.
30. Doll KM, Tseng J, Denslow SA, et al. High-grade endometrial cancer: revisiting the impact of tumor size and location on outcomes. *Gynecol Oncol* 2014;132:44-49. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24183734>.
31. McCluggage WG. My approach to the interpretation of endometrial biopsies and curettings. *J Clin Pathol* 2006;59:801-812. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16873562>.

32. McKenney JK, Longacre TA. Low-grade endometrial adenocarcinoma: a diagnostic algorithm for distinguishing atypical endometrial hyperplasia and other benign (and malignant) mimics. *Adv Anat Pathol* 2009;16:1-22. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19098463>.
33. Leitao MM, Jr., Kehoe S, Barakat RR, et al. Comparison of D&C and office endometrial biopsy accuracy in patients with FIGO grade 1 endometrial adenocarcinoma. *Gynecol Oncol* 2009;113:105-108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19167049>.
34. Gimpelson RJ, Rappold HO. A comparative study between panoramic hysteroscopy with directed biopsies and dilatation and curettage. A review of 276 cases. *Am J Obstet Gynecol* 1988;158:489-492. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3348309>.
35. Lee JH, Dubinsky T, Andreotti RF, et al. ACR appropriateness Criteria(R) pretreatment evaluation and follow-up of endometrial cancer of the uterus. *Ultrasound Q* 2011;27:139-145. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21606818>.
36. Ortashi O, Jain S, Emmanuel O, et al. Evaluation of the sensitivity, specificity, positive and negative predictive values of preoperative magnetic resonance imaging for staging endometrial cancer. A prospective study of 100 cases at the Dorset Cancer Centre. *Eur J Obstet Gynecol Reprod Biol* 2008;137:232-235. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17537566>.
37. Crivellaro C, Signorelli M, Guerra L, et al. Tailoring systematic lymphadenectomy in high-risk clinical early stage endometrial cancer: the role of 18F-FDG PET/CT. *Gynecol Oncol* 2013;130:306-311. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23707673>.
38. Kitajima K, Suzuki K, Senda M, et al. Preoperative nodal staging of uterine cancer: is contrast-enhanced PET/CT more accurate than non-enhanced PET/CT or enhanced CT alone? *Ann Nucl Med* 2011;25:511-519. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21670955>.
39. Antonsen SL, Jensen LN, Loft A, et al. MRI, PET/CT and ultrasound in the preoperative staging of endometrial cancer - a multicenter prospective comparative study. *Gynecol Oncol* 2013;128:300-308. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23200916>.
40. Bollineni VR, Ytre-Hauge S, Bollineni-Balabay O, et al. High Diagnostic Value of 18F-FDG PET/CT in Endometrial Cancer: Systematic Review and Meta-Analysis of the Literature. *J Nucl Med* 2016;57:879-885. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26823564>.
41. Duk JM, Aalders JG, Fleuren GJ, de Bruijn HW. CA 125: a useful marker in endometrial carcinoma. *Am J Obstet Gynecol* 1986;155:1097-1102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3465243>.
42. Duk JM, Aalders JG, Fleuren GJ, et al. Tumor markers CA 125, squamous cell carcinoma antigen, and carcinoembryonic antigen in patients with adenocarcinoma of the uterine cervix. *Obstet Gynecol* 1989;73:661-668. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2648225>.
43. Patsner B, Orr JW, Jr., Mann WJ, Jr. Use of serum CA 125 measurement in posttreatment surveillance of early-stage endometrial carcinoma. *Am J Obstet Gynecol* 1990;162:427-429. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2309825>.
44. Rose PG, Sommers RM, Reale FR, et al. Serial serum CA 125 measurements for evaluation of recurrence in patients with endometrial carcinoma. *Obstet Gynecol* 1994;84:12-16. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8008305>.
45. Price FV, Chambers SK, Carcangiu ML, et al. CA 125 may not reflect disease status in patients with uterine serous carcinoma. *Cancer* 1998;82:1720-1725. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9576294>.
46. Leslie KK, Thiel KW, Goodheart MJ, et al. Endometrial cancer. *Obstet Gynecol Clin North Am* 2012;39:255-268. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22640714>.
47. Smith RA, Brooks D, Cokkinides V, et al. Cancer screening in the United States, 2013: a review of current American Cancer Society

guidelines, current issues in cancer screening, and new guidance on cervical cancer screening and lung cancer screening. *CA Cancer J Clin* 2013;63:88-105. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23378235>.

48. Boronow RC, Morrow CP, Creasman WT, et al. Surgical staging in endometrial cancer: clinical-pathologic findings of a prospective study. *Obstet Gynecol* 1984;63:825-832. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/6728365>.

49. Cowles TA, Magrina JF, Masterson BJ, Capen CV. Comparison of clinical and surgical-staging in patients with endometrial carcinoma. *Obstet Gynecol* 1985;66:413-416. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/4022500>.

50. Creasman WT, Morrow CP, Bundy BN, et al. Surgical pathologic spread patterns of endometrial cancer. A Gynecologic Oncology Group Study. *Cancer* 1987;60:2035-2041. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/3652025>.

51. Benedet JL, Bender H, Jones H, et al. FIGO staging classifications and clinical practice guidelines in the management of gynecologic cancers. FIGO Committee on Gynecologic Oncology. *Int J Gynaecol Obstet* 2000;70:209-262. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11041682>.

52. Wright JD, Barrena Medel NI, Sehoul J, et al. Contemporary management of endometrial cancer. *Lancet* 2012;379:1352-1360. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22444602>.

53. Pecorelli S. Revised FIGO staging for carcinoma of the vulva, cervix, and endometrium. *Int J Gynaecol Obstet* 2009;105:103-104. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19367689>.

54. Creasman W. Revised FIGO staging for carcinoma of the endometrium. *Int J Gynaecol Obstet* 2009;105:109. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19345353>.

55. Mariani A, Dowdy SC, Podratz KC. New surgical staging of endometrial cancer: 20 years later. *Int J Gynaecol Obstet* 2009;105:110-111. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19285672>.

56. Amin MB, Edge SB, Greene FL, et al. *AJCC Cancer Staging Manual*, 8th edition. New York: Springer; 2017.

57. Wethington SL, Barrena Medel NI, Wright JD, Herzog TJ. Prognostic significance and treatment implications of positive peritoneal cytology in endometrial adenocarcinoma: Unraveling a mystery. *Gynecol Oncol* 2009;115:18-25. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19632708>.

58. Takeshima N, Nishida H, Tabata T, et al. Positive peritoneal cytology in endometrial cancer: enhancement of other prognostic indicators. *Gynecol Oncol* 2001;82:470-473. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11520142>.

59. Goff BA, Rice LW. Assessment of depth of myometrial invasion in endometrial adenocarcinoma. *Gynecol Oncol* 1990;38:46-48. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2354826>.

60. Daniel AG, Peters WA, 3rd. Accuracy of office and operating room curettage in the grading of endometrial carcinoma. *Obstet Gynecol* 1988;71:612-614. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/3353053>.

61. Soliman PT, Frumovitz M, Spannuth W, et al. Lymphadenectomy during endometrial cancer staging: practice patterns among gynecologic oncologists. *Gynecol Oncol* 2010;119:291-294. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20708226>.

62. Kumar S, Mariani A, Bakkum-Gamez JN, et al. Risk factors that mitigate the role of paraaortic lymphadenectomy in uterine endometrioid cancer. *Gynecol Oncol* 2013;130:441-445. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23747331>.

63. Kilgore LC, Partridge EE, Alvarez RD, et al. Adenocarcinoma of the endometrium: survival comparisons of patients with and without pelvic

node sampling. *Gynecol Oncol* 1995;56:29-33. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7821843>.

64. Havrilesky LJ, Cragun JM, Calingaert B, et al. Resection of lymph node metastases influences survival in stage IIIC endometrial cancer. *Gynecol Oncol* 2005;99:689-695. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16126261>.

65. Todo Y, Kato H, Kaneuchi M, et al. Survival effect of para-aortic lymphadenectomy in endometrial cancer (SEPAL study): a retrospective cohort analysis. *Lancet* 2010;375:1165-1172. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20188410>.

66. Kitchener H, Swart AM, Qian Q, et al. Efficacy of systematic pelvic lymphadenectomy in endometrial cancer (MRC ASTEC trial): a randomised study. *Lancet* 2009;373:125-136. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19070889>.

67. Benedetti Panici P, Basile S, Maneschi F, et al. Systematic pelvic lymphadenectomy vs. no lymphadenectomy in early-stage endometrial carcinoma: randomized clinical trial. *J Natl Cancer Inst* 2008;100:1707-1716. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19033573>.

68. ACOG practice bulletin, clinical management guidelines for obstetrician-gynecologists, number 65, August 2005: management of endometrial cancer. *Obstet Gynecol* 2005;106:413-425. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16055605>.

69. Seamon LG, Fowler JM, Cohn DE. Lymphadenectomy for endometrial cancer: the controversy. *Gynecol Oncol* 2010;117:6-8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20110120>.

70. Creasman WT, Mutch DE, Herzog TJ. ASTEC lymphadenectomy and radiation therapy studies: are conclusions valid? *Gynecol Oncol* 2010;116:293-294. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19897230>.

71. Uccella S, Podratz KC, Aletti GD, Mariani A. Lymphadenectomy in endometrial cancer. *Lancet* 2009;373:1170; author reply 1170-1171. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19345823>.

72. Uccella S, Podratz KC, Aletti GD, Mariani A. Re: Systematic pelvic lymphadenectomy vs no lymphadenectomy in early-stage endometrial carcinoma: randomized clinical trial. *J Natl Cancer Inst* 2009;101:897-898; author reply 898-899. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19509367>.

73. Milam MR, Java J, Walker JL, et al. Nodal metastasis risk in endometrioid endometrial cancer. *Obstet Gynecol* 2012;119:286-292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22270280>.

74. Neubauer NL, Lurain JR. The role of lymphadenectomy in surgical staging of endometrial cancer. *Int J Surg Oncol* 2011;2011:814649. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22312525>.

75. Mariani A, Dowdy SC, Cliby WA, et al. Prospective assessment of lymphatic dissemination in endometrial cancer: a paradigm shift in surgical staging. *Gynecol Oncol* 2008;109:11-18. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18304622>.

76. Hirahatake K, Hareyama H, Sakuragi N, et al. A clinical and pathologic study on para-aortic lymph node metastasis in endometrial carcinoma. *J Surg Oncol* 1997;65:82-87. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9209518>.

77. Frederick PJ, Straughn JM, Jr. The role of comprehensive surgical staging in patients with endometrial cancer. *Cancer Control* 2009;16:23-29. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19078926>.

78. Barlin JN, Khoury-Collado F, Kim CH, et al. The importance of applying a sentinel lymph node mapping algorithm in endometrial cancer staging: beyond removal of blue nodes. *Gynecol Oncol* 2012;125:531-535. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22366409>.

79. Ballester M, Koskas M, Coutant C, et al. Does the use of the 2009 FIGO classification of endometrial cancer impact on indications of the

sentinel node biopsy? BMC Cancer 2010;10:465. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20804553>.

80. How J, Lau S, Press J, et al. Accuracy of sentinel lymph node detection following intra-operative cervical injection for endometrial cancer: a prospective study. Gynecol Oncol 2012;127:332-337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22910695>.

81. Khoury-Collado F, Murray MP, Hensley ML, et al. Sentinel lymph node mapping for endometrial cancer improves the detection of metastatic disease to regional lymph nodes. Gynecol Oncol 2011;122:251-254. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21570109>.

82. Holloway RW, Abu-Rustum NR, Backes FJ, et al. Sentinel lymph node mapping and staging in endometrial cancer: A Society of Gynecologic Oncology literature review with consensus recommendations. Gynecol Oncol 2017;146:405-415. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28566221>.

83. Vidal F, Leguevaque P, Motton S, et al. Evaluation of the sentinel lymph node algorithm with blue dye labeling for early-stage endometrial cancer in a multicentric setting. Int J Gynecol Cancer 2013;23:1237-1243. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23839245>.

84. Kim CH, Khoury-Collado F, Barber EL, et al. Sentinel lymph node mapping with pathologic ultrastaging: A valuable tool for assessing nodal metastasis in low-grade endometrial cancer with superficial myoinvasion. Gynecol Oncol 2013. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24099838>.

85. Group SGOCPECW, Burke WM, Orr J, et al. Endometrial cancer: a review and current management strategies: part I. Gynecol Oncol 2014;134:385-392. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24905773>.

86. Soliman PT, Westin SN, Dioun S, et al. A prospective validation study of sentinel lymph node mapping for high-risk endometrial cancer. Gynecol Oncol 2017;146:234-239. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28528918>.

87. Abu-Rustum NR, Khoury-Collado F, Pandit-Taskar N, et al. Sentinel lymph node mapping for grade 1 endometrial cancer: is it the answer to the surgical staging dilemma? Gynecol Oncol 2009;113:163-169. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19232699>.

88. Ballester M, Dubernard G, Lecuru F, et al. Detection rate and diagnostic accuracy of sentinel-node biopsy in early stage endometrial cancer: a prospective multicentre study (SENTI-ENDO). Lancet Oncol 2011;12:469-476. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21489874>.

89. Press JZ, Gotlieb WH. Controversies in the treatment of early stage endometrial carcinoma. Obstet Gynecol Int 2012;2012:578490. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22685466>.

90. Touhami O, Trinh XB, Gregoire J, et al. Predictors of non-sentinel lymph node (non-SLN) metastasis in patients with sentinel lymph node (SLN) metastasis in endometrial cancer. Gynecol Oncol 2015;138:41-45. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25891803>.

91. Rossi EC, Kowalski LD, Scalici J, et al. A comparison of sentinel lymph node biopsy to lymphadenectomy for endometrial cancer staging (FIRES trial): a multicentre, prospective, cohort study. Lancet Oncol 2017;18:384-392. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28159465>.

92. Paley PJ, Veljovich DS, Press JZ, et al. A prospective investigation of fluorescence imaging to detect sentinel lymph nodes at robotic-assisted endometrial cancer staging. Am J Obstet Gynecol 2016;215:117 e111-117. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26743505>.

93. Sinno AK, Fader AN, Roche KL, et al. A comparison of colorimetric versus fluorometric sentinel lymph node mapping during robotic surgery for endometrial cancer. Gynecol Oncol 2014;134:281-286. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24882555>.

94. Ruscito I, Gasparri ML, Braicu EI, et al. Sentinel Node Mapping in Cervical and Endometrial Cancer: Indocyanine Green Versus Other Conventional Dyes-A Meta-Analysis. Ann Surg Oncol 2016;23:3749-3756. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27160526>.

95. Bodurtha Smith AJ, Fader AN, Tanner EJ. Sentinel lymph node assessment in endometrial cancer: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2016. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27871836>.

96. Cormier B, Rozenholc AT, Gotlieb W, et al. Sentinel lymph node procedure in endometrial cancer: A systematic review and proposal for standardization of future research. *Gynecol Oncol* 2015;138:478-485. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26047592>.

97. Naoura I, Canlorbe G, Bendifallah S, et al. Relevance of sentinel lymph node procedure for patients with high-risk endometrial cancer. *Gynecol Oncol* 2015;136:60-64. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25449312>.

98. Schiavone MB, Zivanovic O, Zhou Q, et al. Survival of Patients with Uterine Carcinosarcoma Undergoing Sentinel Lymph Node Mapping. *Ann Surg Oncol* 2016;23:196-202. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25994210>.

99. Holloway RW, Gupta S, Stavitzski NM, et al. Sentinel lymph node mapping with staging lymphadenectomy for patients with endometrial cancer increases the detection of metastasis. *Gynecol Oncol* 2016;141:206-210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26905211>.

100. Darai E, Dubernard G, Bats AS, et al. Sentinel node biopsy for the management of early stage endometrial cancer: long-term results of the SENTI-ENDO study. *Gynecol Oncol* 2015;136:54-59. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25450151>.

101. Kim CH, Soslow RA, Park KJ, et al. Pathologic ultrastaging improves micrometastasis detection in sentinel lymph nodes during endometrial cancer staging. *Int J Gynecol Cancer* 2013;23:964-970. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23694985>.

102. Raimond E, Ballester M, Hudry D, et al. Impact of sentinel lymph node biopsy on the therapeutic management of early-stage endometrial cancer: Results of a retrospective multicenter study. *Gynecol Oncol*

2014;133:506-511. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24642092>.

103. Frimer M, Khoury-Collado F, Murray MP, et al. Micrometastasis of endometrial cancer to sentinel lymph nodes: is it an artifact of uterine manipulation? *Gynecol Oncol* 2010;119:496-499. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20888626>.

104. Touboul C, Bentivegna E, Uzan C, et al. Sentinel lymph node in endometrial cancer: a review. *Curr Oncol Rep* 2013;15:559-565. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24190831>.

105. Amezcua CA, MacDonald HR, Lum CA, et al. Endometrial cancer patients have a significant risk of harboring isolated tumor cells in histologically negative lymph nodes. *Int J Gynecol Cancer* 2006;16:1336-1341. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16803526>.

106. Todo Y, Kato H, Okamoto K, et al. Isolated tumor cells and micrometastases in regional lymph nodes in stage I to II endometrial cancer. *J Gynecol Oncol* 2016;27:e1. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25925293>.

107. Smerage JB, Barlow WE, Hortobagyi GN, et al. Circulating tumor cells and response to chemotherapy in metastatic breast cancer: SWOG S0500. *J Clin Oncol* 2014;32:3483-3489. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24888818>.

108. St Clair CM, Eriksson AG, Ducie JA, et al. Low-Volume Lymph Node Metastasis Discovered During Sentinel Lymph Node Mapping for Endometrial Carcinoma. *Ann Surg Oncol* 2016;23:1653-1659. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26714954>.

109. Plante M, Stanleigh J, Renaud MC, et al. Isolated tumor cells identified by sentinel lymph node mapping in endometrial cancer: Does adjuvant treatment matter? *Gynecol Oncol* 2017;146:240-246. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28577885>.

110. Scalici J, Laughlin BB, Finan MA, et al. The trend towards minimally invasive surgery (MIS) for endometrial cancer: an ACS-NSQIP evaluation

of surgical outcomes. *Gynecol Oncol* 2015;136:512-515. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25462206>.

111. Janda M, GebSKI V, Davies LC, et al. Effect of Total Laparoscopic Hysterectomy vs Total Abdominal Hysterectomy on Disease-Free Survival Among Women With Stage I Endometrial Cancer: A Randomized Clinical Trial. *JAMA* 2017;317:1224-1233. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28350928>.

112. Fader AN, Weise RM, Sinno AK, et al. Utilization of Minimally Invasive Surgery in Endometrial Cancer Care: A Quality and Cost Disparity. *Obstet Gynecol* 2016;127:91-100. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26646127>.

113. Galaal K, Bryant A, Fisher AD, et al. Laparoscopy versus laparotomy for the management of early stage endometrial cancer. *Cochrane Database Syst Rev* 2012:CD006655. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22972096>.

114. Kornblith AB, Huang HQ, Walker JL, et al. Quality of life of patients with endometrial cancer undergoing laparoscopic international federation of gynecology and obstetrics staging compared with laparotomy: a Gynecologic Oncology Group study. *J Clin Oncol* 2009;27:5337-5342. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19805678>.

115. Walker JL, Piedmonte MR, Spirtos NM, et al. Laparoscopy compared with laparotomy for comprehensive surgical staging of uterine cancer: Gynecologic Oncology Group Study LAP2. *J Clin Oncol* 2009;27:5331-5336. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19805679>.

116. Mannschreck D, Matsuno RK, Moriarty JP, et al. Disparities in Surgical Care Among Women With Endometrial Cancer. *Obstet Gynecol* 2016;128:526-534. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27500330>.

117. Walker JL, Piedmonte MR, Spirtos NM, et al. Recurrence and survival after random assignment to laparoscopy versus laparotomy for comprehensive surgical staging of uterine cancer: Gynecologic Oncology

Group LAP2 Study. *J Clin Oncol* 2012;30:695-700. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22291074>.

118. King LP, Miller DS. Recent progress: gynecologic oncology group trials in uterine corpus tumors. *Rev Recent Clin Trials* 2009;4:70-74. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19463102>.

119. Vergote I, Amant F, Neven P. Is it safe to treat endometrial carcinoma endoscopically? *J Clin Oncol* 2009;27:5305-5307. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19805666>.

120. Mourits MJ, Bijen CB, Arts HJ, et al. Safety of laparoscopy versus laparotomy in early-stage endometrial cancer: a randomised trial. *Lancet Oncol* 2010;11:763-771. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20638901>.

121. He H, Zeng D, Ou H, et al. Laparoscopic treatment of endometrial cancer: systematic review. *J Minim Invasive Gynecol* 2013;20:413-423. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23506718>.

122. Wang HL, Ren YF, Yang J, et al. Total laparoscopic hysterectomy versus total abdominal hysterectomy for endometrial cancer: a meta-analysis. *Asian Pac J Cancer Prev* 2013;14:2515-2519. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23725166>.

123. Mori KM, Neubauer NL. Minimally invasive surgery in gynecologic oncology. *ISRN Obstet Gynecol* 2013;2013:312982. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23997959>.

124. Krill LS, Bristow RE. Robotic surgery: gynecologic oncology. *Cancer J* 2013;19:167-176. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23528726>.

125. ElSahwi KS, Hooper C, De Leon MC, et al. Comparison between 155 cases of robotic vs. 150 cases of open surgical staging for endometrial cancer. *Gynecol Oncol* 2012;124:260-264. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22036203>.

126. Chan JK, Gardner AB, Taylor K, et al. Robotic versus laparoscopic versus open surgery in morbidly obese endometrial cancer patients - a comparative analysis of total charges and complication rates. *Gynecol Oncol* 2015;139:300-305. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26363212>.
127. Coronado PJ, Herraiz MA, Magrina JF, et al. Comparison of perioperative outcomes and cost of robotic-assisted laparoscopy, laparoscopy and laparotomy for endometrial cancer. *Eur J Obstet Gynecol Reprod Biol* 2012;165:289-294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22819573>.
128. Seamon LG, Cohn DE, Henretta MS, et al. Minimally invasive comprehensive surgical staging for endometrial cancer: Robotics or laparoscopy? *Gynecol Oncol* 2009;113:36-41. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19168206>.
129. Bell MC, Torgerson J, Seshadri-Kreaden U, et al. Comparison of outcomes and cost for endometrial cancer staging via traditional laparotomy, standard laparoscopy and robotic techniques. *Gynecol Oncol* 2008;111:407-411. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18829091>.
130. Cardenas-Goicoechea J, Adams S, Bhat SB, Randall TC. Surgical outcomes of robotic-assisted surgical staging for endometrial cancer are equivalent to traditional laparoscopic staging at a minimally invasive surgical center. *Gynecol Oncol* 2010;117:224-228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20144471>.
131. Brudie LA, Backes FJ, Ahmad S, et al. Analysis of disease recurrence and survival for women with uterine malignancies undergoing robotic surgery. *Gynecol Oncol* 2013;128:309-315. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23153590>.
132. Backes FJ, Brudie LA, Farrell MR, et al. Short- and long-term morbidity and outcomes after robotic surgery for comprehensive endometrial cancer staging. *Gynecol Oncol* 2012;125:546-551. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22387522>.
133. Fleming ND, Ramirez PT. Robotic surgery in gynecologic oncology. *Curr Opin Oncol* 2012;24:547-553. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22581356>.
134. Siesto G, Ornaghi S, Ieda N, Vitobello D. Robotic surgical staging for endometrial and cervical cancers in medically ill patients. *Gynecol Oncol* 2013;129:593-597. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23454499>.
135. van Dam P, Hauspy J, Verkinderen L, et al. Are costs of robot-assisted surgery warranted for gynecological procedures? *Obstet Gynecol Int* 2011;2011:973830. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21941556>.
136. Weinberg L, Rao S, Escobar PF. Robotic surgery in gynecology: an updated systematic review. *Obstet Gynecol Int* 2011;2011:852061. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22190948>.
137. Ramirez PT, Adams S, Boggess JF, et al. Robotic-assisted surgery in gynecologic oncology: a Society of Gynecologic Oncology consensus statement. Developed by the Society of Gynecologic Oncology's Clinical Practice Robotics Task Force. *Gynecol Oncol* 2012;124:180-184. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22079679>.
138. AAGL. Guidelines for privileging for robotic-assisted gynecologic laparoscopy. *J Minim Invasive Gynecol* 2014;21:157-167. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24509290>.
139. American Congress of Obstetricians and Gynecologists. Statement on Robotic Surgery by ACOG President James T. Breedon. 2013. Available at: <http://www.acog.org/About-ACOG/News-Room/News-Releases/2013/Statement-on-Robotic-Surgery>. Accessed
140. Sinno AK, Fader AN. Robotic-assisted surgery in gynecologic oncology. *Fertil Steril* 2014;102:922-932. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25274485>.
141. Gala RB, Margulies R, Steinberg A, et al. Systematic review of robotic surgery in gynecology: robotic techniques compared with

laparoscopy and laparotomy. J Minim Invasive Gynecol 2014;21:353-361. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24295923>.

142. SGO Position Statement: Morcellation. Society of Gynecologic Oncology; 2013. Available at: <https://www.sgo.org/newsroom/position-statements-2/morcellation/>. Accessed September 30, 2014.

143. Power Morcellation and Occult Malignancy in Gynecologic Surgery. The American College of Obstetrics and Gynecologists; 2014. Available at: <http://www.acog.org/Resources-And-Publications/Task-Force-and-Work-Group-Reports/Power-Morcellation-and-Occult-Malignancy-in-Gynecologic-Surgery>. Accessed September 30, 2014.

144. U.S. Department of Health and Human Services. FDA discourages use of laparoscopic power morcellation for removal of uterus or uterine fibroids Food and Drug Administration; 2014. Available at: <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm393689.htm>. Accessed September 30, 2014.

145. Bogani G, Cliby WA, Aletti GD. Impact of morcellation on survival outcomes of patients with unexpected uterine leiomyosarcoma: a systematic review and meta-analysis. Gynecol Oncol 2015;137:167-172. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25462199>.

146. Wright JD, Buck AM, Shah M, et al. Safety of ovarian preservation in premenopausal women with endometrial cancer. J Clin Oncol 2009;27:1214-1219. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19171707>.

147. Koskas M, Bendifallah S, Luton D, et al. Safety of uterine and/or ovarian preservation in young women with grade 1 intramucous endometrial adenocarcinoma: a comparison of survival according to the extent of surgery. Fertil Steril 2012;98:1229-1235. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22959452>.

148. Lee TS, Lee JY, Kim JW, et al. Outcomes of ovarian preservation in a cohort of premenopausal women with early-stage endometrial cancer: A Korean Gynecologic Oncology Group study. Gynecol Oncol

2013;131:289-293. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23994534>.

149. Manfredi R, Mirk P, Maresca G, et al. Local-regional staging of endometrial carcinoma: role of MR imaging in surgical planning. Radiology 2004;231:372-378. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15031434>.

150. Akin O, Mironov S, Pandit-Taskar N, Hann LE. Imaging of uterine cancer. Radiol Clin North Am 2007;45:167-182. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17157628>.

151. Gunderson CC, Fader AN, Carson KA, Bristow RE. Oncologic and reproductive outcomes with progestin therapy in women with endometrial hyperplasia and grade 1 adenocarcinoma: a systematic review. Gynecol Oncol 2012;125:477-482. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22245711>.

152. Baker J, Obermair A, GebSKI V, Janda M. Efficacy of oral or intrauterine device-delivered progestin in patients with complex endometrial hyperplasia with atypia or early endometrial adenocarcinoma: a meta-analysis and systematic review of the literature. Gynecol Oncol 2012;125:263-270. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22196499>.

153. Gracia CR, Jeruss JS. Lives in the balance: women with cancer and the right to fertility care. J Clin Oncol 2013;31:668-669. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23341520>.

154. Ushijima K, Yahata H, Yoshikawa H, et al. Multicenter phase II study of fertility-sparing treatment with medroxyprogesterone acetate for endometrial carcinoma and atypical hyperplasia in young women. J Clin Oncol 2007;25:2798-2803. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17602085>.

155. Hubbs JL, Saig RM, Abaid LN, et al. Systemic and local hormone therapy for endometrial hyperplasia and early adenocarcinoma. Obstet Gynecol 2013;121:1172-1180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23812449>.

156. Trimble CL, Method M, Leitao M, et al. Management of endometrial precancers. *Obstet Gynecol* 2012;120:1160-1175. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23090535>.

157. Mehaseb MK, Latimer JA. Controversies in the management of endometrial carcinoma: an update. *Obstet Gynecol Int* 2012;2012:676032. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22518164>.

158. Hahn HS, Yoon SG, Hong JS, et al. Conservative treatment with progestin and pregnancy outcomes in endometrial cancer. *Int J Gynecol Cancer* 2009;19:1068-1073. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19820370>.

159. Park JY, Kim DY, Kim JH, et al. Long-term oncologic outcomes after fertility-sparing management using oral progestin for young women with endometrial cancer (KGOG 2002). *Eur J Cancer* 2013;49:868-874. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23072814>.

160. Park JY, Seong SJ, Kim TJ, et al. Pregnancy outcomes after fertility-sparing management in young women with early endometrial cancer. *Obstet Gynecol* 2013;121:136-142. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23262938>.

161. Boente MP, Yordan EL, Jr., McIntosh DG, et al. Prognostic factors and long-term survival in endometrial adenocarcinoma with cervical involvement. *Gynecol Oncol* 1993;51:316-322. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8112639>.

162. Sartori E, Gadducci A, Landoni F, et al. Clinical behavior of 203 stage II endometrial cancer cases: the impact of primary surgical approach and of adjuvant radiation therapy. *Int J Gynecol Cancer* 2001;11:430-437. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11906545>.

163. Gadducci A, Cosio S, Genazzani AR. Old and new perspectives in the pharmacological treatment of advanced or recurrent endometrial cancer: Hormonal therapy, chemotherapy and molecularly targeted therapies. *Crit Rev Oncol Hematol* 2006;58:242-256. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16436330>.

164. Mountzios G, Pectasides D, Bournakis E, et al. Developments in the systemic treatment of endometrial cancer. *Crit Rev Oncol Hematol* 2011;79:278-292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20833559>.

165. Fiorica JV, Brunetto VL, Hanjani P, et al. Phase II trial of alternating courses of megestrol acetate and tamoxifen in advanced endometrial carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol* 2004;92:10-14. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14751131>.

166. Altman AD, Thompson J, Nelson G, et al. Use of aromatase inhibitors as first- and second-line medical therapy in patients with endometrial adenocarcinoma: a retrospective study. *J Obstet Gynaecol Can* 2012;34:664-672. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22742486>.

167. Barker LC, Brand IR, Crawford SM. Sustained effect of the aromatase inhibitors anastrozole and letrozole on endometrial thickness in patients with endometrial hyperplasia and endometrial carcinoma. *Curr Med Res Opin* 2009;25:1105-1109. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19301987>.

168. Rose PG, Brunetto VL, VanLe L, et al. A phase II trial of anastrozole in advanced recurrent or persistent endometrial carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol* 2000;78:212-216. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10926805>.

169. Decruze SB, Green JA. Hormone therapy in advanced and recurrent endometrial cancer: a systematic review. *Int J Gynecol Cancer* 2007;17:964-978. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17442022>.

170. Fishman DA, Roberts KB, Chambers JT, et al. Radiation therapy as exclusive treatment for medically inoperable patients with stage I and II endometrioid carcinoma with endometrium. *Gynecol Oncol* 1996;61:189-196. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8626131>.

171. Coon D, Beriwal S, Heron DE, et al. High-dose-rate Rotte "Y" applicator brachytherapy for definitive treatment of medically inoperable endometrial cancer: 10-year results. *Int J Radiat Oncol Biol Phys* 2008;71:779-783. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18258388>.

172. Niazi TM, Souhami L, Portelance L, et al. Long-term results of high-dose-rate brachytherapy in the primary treatment of medically inoperable stage I-II endometrial carcinoma. *Int J Radiat Oncol Biol Phys* 2005;63:1108-1113. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16099598>.

173. van der Steen-Banasik E, Christiaens M, Shash E, et al. Systemic review: Radiation therapy alone in medical non-operable endometrial carcinoma. *Eur J Cancer* 2016;65:172-181. Available at:

174. Barlin JN, Puri I, Bristow RE. Cytoreductive surgery for advanced or recurrent endometrial cancer: a meta-analysis. *Gynecol Oncol* 2010;118:14-18. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20434198>.

175. Landrum LM, Moore KN, Myers TK, et al. Stage IVB endometrial cancer: does applying an ovarian cancer treatment paradigm result in similar outcomes? A case-control analysis. *Gynecol Oncol* 2009;112:337-341. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19041126>.

176. Lambrou NC, Gomez-Marin O, Mirhashemi R, et al. Optimal surgical cytoreduction in patients with Stage III and Stage IV endometrial carcinoma: a study of morbidity and survival. *Gynecol Oncol* 2004;93:653-658. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15196860>.

177. Smith SC, Koh WJ. Palliative radiation therapy for gynaecological malignancies. *Best Pract Res Clin Obstet Gynaecol* 2001;15:265-278. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11358401>.

178. Randall ME, Filiaci VL, Muss H, et al. Randomized phase III trial of whole-abdominal irradiation versus doxorubicin and cisplatin chemotherapy in advanced endometrial carcinoma: a Gynecologic

Oncology Group Study. *J Clin Oncol* 2006;24:36-44. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16330675>.

179. Homesley HD, Filiaci V, Gibbons SK, et al. A randomized phase III trial in advanced endometrial carcinoma of surgery and volume directed radiation followed by cisplatin and doxorubicin with or without paclitaxel: A Gynecologic Oncology Group study. *Gynecol Oncol* 2009;112:543-552. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19108877>.

180. Secord AA, Havrilesky LJ, O'Malley DM, et al. A multicenter evaluation of sequential multimodality therapy and clinical outcome for the treatment of advanced endometrial cancer. *Gynecol Oncol* 2009;114:442-447. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19560193>.

181. Creutzberg CL, van Stiphout RG, Nout RA, et al. Nomograms for prediction of outcome with or without adjuvant radiation therapy for patients with endometrial cancer: a pooled analysis of PORTEC-1 and PORTEC-2 trials. *Int J Radiat Oncol Biol Phys* 2015;91:530-539. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25680597>.

182. Group SGOPECW, Burke WM, Orr J, et al. Endometrial cancer: a review and current management strategies: part II. *Gynecol Oncol* 2014;134:393-402. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24929052>.

183. Creutzberg CL, Nout RA. The role of radiotherapy in endometrial cancer: current evidence and trends. *Curr Oncol Rep* 2011;13:472-478. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21845420>.

184. Klopp A, Smith BD, Alektiar K, et al. The role of postoperative radiation therapy for endometrial cancer: Executive summary of an American Society for Radiation Oncology evidence-based guideline. *Pract Radiat Oncol* 2014;4:137-144. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24766678>.

185. Meyer LA, Bohlke K, Powell MA, et al. Postoperative Radiation Therapy for Endometrial Cancer: American Society of Clinical Oncology Clinical Practice Guideline Endorsement of the American Society for Radiation Oncology Evidence-Based Guideline. *J Clin Oncol*

2015;33:2908-2913. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26150442>.

186. Morrow CP, Bundy BN, Kurman RJ, et al. Relationship between surgical-pathological risk factors and outcome in clinical stage I and II carcinoma of the endometrium: a Gynecologic Oncology Group study. *Gynecol Oncol* 1991;40:55-65. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/1989916>.

187. Neubauer NL, Havrilesky LJ, Calingaert B, et al. The role of lymphadenectomy in the management of preoperative grade 1 endometrial carcinoma. *Gynecol Oncol* 2009;112:511-516. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19144394>.

188. Gretz HFr, Economos K, Husain A, et al. The practice of surgical staging and its impact on adjuvant treatment recommendations in patients with stage I endometrial carcinoma. *Gynecol Oncol* 1996;61:409-415. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8641624>.

189. Ben-Shachar I, Pavelka J, Cohn DE, et al. Surgical staging for patients presenting with grade 1 endometrial carcinoma. *Obstet Gynecol* 2005;105:487-493. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15738013>.

190. Elshaikh MA, Al-Wahab Z, Mahdi H, et al. Recurrence patterns and survival endpoints in women with stage II uterine endometrioid carcinoma: a multi-institution study. *Gynecol Oncol* 2015;136:235-239. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25511158>.

191. Creutzberg CL, van Putten WL, Koper PC, et al. Surgery and postoperative radiotherapy versus surgery alone for patients with stage-1 endometrial carcinoma: multicentre randomised trial. PORTEC Study Group. *Post Operative Radiation Therapy in Endometrial Carcinoma*. *Lancet* 2000;355:1404-1411. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10791524>.

192. Aalders J, Abeler V, Kolstad P, Onsrud M. Postoperative external irradiation and prognostic parameters in stage I endometrial carcinoma: clinical and histopathologic study of 540 patients. *Obstet Gynecol*

1980;56:419-427. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/6999399>.

193. Blake P, Swart AM, Orton J, et al. Adjuvant external beam radiotherapy in the treatment of endometrial cancer (MRC ASTEC and NCIC CTG EN.5 randomised trials): pooled trial results, systematic review, and meta-analysis. *Lancet* 2009;373:137-146. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19070891>.

194. Keys HM, Roberts JA, Brunetto VL, et al. A phase III trial of surgery with or without adjunctive external pelvic radiation therapy in intermediate risk endometrial adenocarcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol* 2004;92:744-751. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/14984936>.

195. Creutzberg CL, van Putten WLJ, Wárlám-Rodenhuis CC, et al. Outcome of high-risk stage IC, grade 3, compared with stage I endometrial carcinoma patients: the Postoperative Radiation Therapy in Endometrial Carcinoma Trial. *J Clin Oncol* 2004;22:1234-1241. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15051771>.

196. Scholten AN, van Putten WLJ, Beerman H, et al. Postoperative radiotherapy for stage 1 endometrial carcinoma: long-term outcome of the randomized PORTEC trial with central pathology review. *Int J Radiat Oncol Biol Phys* 2005;63:834-838. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15927414>.

197. Hockel M, Dornhofer N. Treatment of early endometrial carcinoma: is less more? *Lancet* 2009;373:97-99. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19070890>.

198. Creutzberg CL, Nout RA, Lybeert ML, et al. Fifteen-year radiotherapy outcomes of the randomized PORTEC-1 trial for endometrial carcinoma. *Int J Radiat Oncol Biol Phys* 2011;81:e631-638. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21640520>.

199. Alektiar KM, Venkatraman E, Chi DS, Barakat RR. Intravaginal brachytherapy alone for intermediate-risk endometrial cancer. *Int J Radiat*

Oncol Biol Phys 2005;62:111-117. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/15850910>.

200. Nout RA, Smit VT, Putter H, et al. Vaginal brachytherapy versus pelvic external beam radiotherapy for patients with endometrial cancer of high-intermediate risk (PORTEC-2): an open-label, non-inferiority, randomised trial. *Lancet* 2010;375:816-823. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/20206777>.

201. Small W, Jr., Beriwal S, Demanes DJ, et al. American Brachytherapy Society consensus guidelines for adjuvant vaginal cuff brachytherapy after hysterectomy. *Brachytherapy* 2012;11:58-67. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/22265439>.

202. Onsrud M, Cvancarova M, Hellebust TP, et al. Long-term outcomes after pelvic radiation for early-stage endometrial cancer. *J Clin Oncol* 2013;31:3951-3956. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/24019546>.

203. Nout RA, Putter H, Jurgenliemk-Schulz IM, et al. Five-year quality of life of endometrial cancer patients treated in the randomised Post Operative Radiation Therapy in Endometrial Cancer (PORTEC-2) trial and comparison with norm data. *Eur J Cancer* 2012;48:1638-1648. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22176868>.

204. Roper B, Astner ST, Heydemann-Obradovic A, et al. Ten-year data on 138 patients with endometrial carcinoma and postoperative vaginal brachytherapy alone: no need for external-beam radiotherapy in low and intermediate risk patients. *Gynecol Oncol* 2007;107:541-548. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/17884152>.

205. Nout RA, Putter H, Jurgenliemk-Schulz IM, et al. Quality of life after pelvic radiotherapy or vaginal brachytherapy for endometrial cancer: first results of the randomized PORTEC-2 trial. *J Clin Oncol* 2009;27:3547-3556. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19546404>.

206. McCloskey SA, Tchabo NE, Malhotra HK, et al. Adjuvant vaginal brachytherapy alone for high risk localized endometrial cancer as defined by the three major randomized trials of adjuvant pelvic radiation. *Gynecol*

Oncol 2010;116:404-407. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/19944453>.

207. Dunn EF, Geye H, Platta CS, et al. Predictive factors of recurrence following adjuvant vaginal cuff brachytherapy alone for stage I endometrial cancer. *Gynecol Oncol* 2014;133:494-498. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/24657301>.

208. Randall ME, Filiaci V, McMeekin DS, et al. A Phase 3 Trial of Pelvic Radiation Therapy Versus Vaginal Cuff Brachytherapy Followed by Paclitaxel/Carboplatin Chemotherapy in Patients with High-Risk, Early-Stage Endometrial Cancer: A Gynecology Oncology Group Study [abstract]. *Int J Radiat Oncol Biol Phys* 2017. Available at:  
[https://www.astro.org/uploadedFiles/MAIN\\_SITE/Meetings\\_and\\_Education/ASTRO\\_Meetings/2017/Annual\\_Meeting/Content\\_Pieces/Late-breakingAbstracts.pdf](https://www.astro.org/uploadedFiles/MAIN_SITE/Meetings_and_Education/ASTRO_Meetings/2017/Annual_Meeting/Content_Pieces/Late-breakingAbstracts.pdf).

209. Chino JP, Jones E, Berchuck A, et al. The Influence of Radiation Modality and Lymph Node Dissection on Survival in Early-stage Endometrial Cancer. *Int J Radiat Oncol Biol Phys* 2011. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/21640502>.

210. Lee CM, Szabo A, Shrieve DC, et al. Frequency and effect of adjuvant radiation therapy among women with stage I endometrial adenocarcinoma. *JAMA* 2006;295:389-397. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/16434629>.

211. Johnson N, Cornes P. Survival and recurrent disease after postoperative radiotherapy for early endometrial cancer: systematic review and meta-analysis. *BJOG* 2007;114:1313-1320. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/17803718>.

212. Kong A, Johnson N, Cornes P, et al. Adjuvant radiotherapy for stage I endometrial cancer. *Cochrane Database Syst Rev* 2007. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/17443533>.

213. Kong A, Johnson N, Kitchener HC, Lawrie TA. Adjuvant radiotherapy for stage I endometrial cancer: an updated Cochrane systematic review

and meta-analysis. *J Natl Cancer Inst* 2012;104:1625-1634. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22962693>.

214. Eifel PJ. The role of adjuvant radiation therapy for stage I endometrial cancer: does meta-analysis reveal the answer? *J Natl Cancer Inst* 2012;104:1615-1616. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23104209>.

215. Park HJ, Nam EJ, Kim S, et al. The benefit of adjuvant chemotherapy combined with postoperative radiotherapy for endometrial cancer: a meta-analysis. *Eur J Obstet Gynecol Reprod Biol* 2013;170:39-44. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23810000>.

216. Hogberg T, Signorelli M, de Oliveira CF, et al. Sequential adjuvant chemotherapy and radiotherapy in endometrial cancer--results from two randomised studies. *Eur J Cancer* 2010;46:2422-2431. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20619634>.

217. Johnson N, Bryant A, Miles T, et al. Adjuvant chemotherapy for endometrial cancer after hysterectomy. *Cochrane Database Syst Rev* 2011:CD003175. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21975736>.

218. Mustea A, Koensgen D, Belau A, et al. Adjuvant sequential chemoradiation therapy in high-risk endometrial cancer: results of a prospective, multicenter phase-II study of the NOGGO (North-Eastern German Society of Gynaecological Oncology). *Cancer Chemother Pharmacol* 2013;72:975-983. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23995698>.

219. Jutzi L, Hoskins P, Lim P, et al. The importance of adjuvant chemotherapy and pelvic radiotherapy in high-risk early stage endometrial carcinoma. *Gynecol Oncol* 2013;131:581-585. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24055614>.

220. de Boer SM, Powell ME, Mileskin L, et al. Toxicity and quality of life after adjuvant chemoradiotherapy versus radiotherapy alone for women with high-risk endometrial cancer (PORTEC-3): an open-label, multicentre,

randomised, phase 3 trial. *Lancet Oncol* 2016;17:1114-1126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27397040>.

221. Hogberg T. Adjuvant chemotherapy in endometrial carcinoma: overview of randomised trials. *Clin Oncol (R Coll Radiol)* 2008;20:463-469. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18467080>.

222. Koh WJ, Tran AB, Douglas JG, Stelzer KJ. Radiation therapy in endometrial cancer. *Best Pract Res Clin Obstet Gynaecol* 2001;15:417-432. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11476563>.

223. Secord AA, Geller MA, Broadwater G, et al. A multicenter evaluation of adjuvant therapy in women with optimally resected stage IIIC endometrial cancer. *Gynecol Oncol* 2013;128:65-70. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23085460>.

224. Greven KM, Lanciano RM, Corn B, et al. Pathologic stage III endometrial carcinoma. Prognostic factors and patterns of recurrence. *Cancer* 1993;71:3697-3702. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8490920>.

225. Gibbons S, Martinez A, Schray M, et al. Adjuvant whole abdominopelvic irradiation for high risk endometrial carcinoma. *Int J Radiat Oncol Biol Phys* 1991;21:1019-1025. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1917597>.

226. Greer BE, Hamberger AD. Treatment of intraperitoneal metastatic adenocarcinoma of the endometrium by the whole-abdomen moving-strip technique and pelvic boost irradiation. *Gynecol Oncol* 1983;16:365-373. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6654180>.

227. Abaid LN, Rettenmaier MA, Brown JV, 3rd, et al. Sequential chemotherapy and radiotherapy as sandwich therapy for the treatment of high risk endometrial cancer. *J Gynecol Oncol* 2012;23:22-27. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22355463>.

228. Geller MA, Ivy JJ, Ghebre R, et al. A phase II trial of carboplatin and docetaxel followed by radiotherapy given in a "Sandwich" method for stage III, IV, and recurrent endometrial cancer. *Gynecol Oncol*



# NCCN Guidelines Version 4.2019

## Uterine Neoplasms

[NCCN Guidelines Index](#)  
[Table of Contents](#)  
[Discussion](#)

2011;121:112-117. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21239048>.

229. Brown AP, Gaffney DK, Dodson MK, et al. Survival analysis of endometrial cancer patients with positive lymph nodes. *Int J Gynecol Cancer* 2013;23:861-868. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23598890>.

230. Lum MM, Belnap TW, Frandsen J, et al. Survival Analysis of Cancer Patients With FIGO Stage IIIA Endometrial Cancer. *Am J Clin Oncol* 2015;38:283-288. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23774072>.

231. Ren Y, Huang X, Shan B, et al. Adjuvant concurrent chemoradiation followed by chemotherapy for high-risk endometrial cancer. *Gynecol Oncol* 2016;140:58-63. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26607778>.

232. Le T, Menard C, Samant R, et al. Longitudinal assessments of quality of life in endometrial cancer patients: effect of surgical approach and adjuvant radiotherapy. *Int J Radiat Oncol Biol Phys* 2009;75:795-802. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19250764>.

233. de Boer SM, Nout RA, Jurgenliemk-Schulz IM, et al. Long-Term Impact of Endometrial Cancer Diagnosis and Treatment on Health-Related Quality of Life and Cancer Survivorship: Results From the Randomized PORTEC-2 Trial. *Int J Radiat Oncol Biol Phys* 2015;93:797-809. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26530748>.

234. Salani R, Khanna N, Frimer M, et al. An update on post-treatment surveillance and diagnosis of recurrence in women with gynecologic malignancies: Society of Gynecologic Oncology (SGO) recommendations. *Gynecol Oncol* 2017;146:3-10. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28372871>.

235. Fung-Kee-Fung M, Dodge J, Elit L, et al. Follow-up after primary therapy for endometrial cancer: a systematic review. *Gynecol Oncol* 2006;101:520-529. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16556457>.

236. Lajer H, Elnegaard S, Christensen RD, et al. Survival after stage IA endometrial cancer; can follow-up be altered? A prospective nationwide Danish survey. *Acta Obstet Gynecol Scand* 2012;91:976-982. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22548255>.

237. Greer BE, Goff BA, Koh WJ. Endometrial carcinoma. In: Johnson FE, Virgo KS, eds. *Cancer Patient Follow-up*. St. Louis: Mosby; 1997:357-377.

238. Bristow RE, Purinton SC, Santillan A, et al. Cost-effectiveness of routine vaginal cytology for endometrial cancer surveillance. *Gynecol Oncol* 2006;103:709-713. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16797686>.

239. Cowens-Alvarado R, Sharpe K, Pratt-Chapman M, et al. Advancing survivorship care through the National Cancer Survivorship Resource Center: developing American Cancer Society guidelines for primary care providers. *CA Cancer J Clin* 2013;63:147-150. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23512728>.

240. McCabe MS, Bhatia S, Oeffinger KC, et al. American Society of Clinical Oncology statement: achieving high-quality cancer survivorship care. *J Clin Oncol* 2013;31:631-640. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23295805>.

241. Smith WA, Nolan VG, Robison LL, et al. Physical activity among cancer survivors and those with no history of cancer- a report from the National Health and Nutrition Examination Survey 2003-2006. *Am J Transl Res* 2011;3:342-350. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21904654>.

242. Cooper AL, Dornfeld-Finke JM, Banks HW, et al. Is cytologic screening an effective surveillance method for detection of vaginal recurrence of uterine cancer? *Obstet Gynecol* 2006;107:71-76. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16394042>.

243. Salani R, Nagel CI, Drennen E, Bristow RE. Recurrence patterns and surveillance for patients with early stage endometrial cancer. *Gynecol Oncol* 2011;123:205-207. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21820709>.

244. Hunn J, Tenney ME, Tergas AI, et al. Patterns and utility of routine surveillance in high grade endometrial cancer. *Gynecol Oncol* 2015;137:485-489. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25838164>.

245. Smith DC, Prentice R, Thompson DJ, Herrmann WL. Association of exogenous estrogen and endometrial carcinoma. *N Engl J Med* 1975;293:1164-1167. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1186789>.

246. Ziel HK, Finkle WD. Increased risk of endometrial carcinoma among users of conjugated estrogens. *N Engl J Med* 1975;293:1167-1170. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/171569>.

247. Creasman WT, Henderson D, Hinshaw W, Clarke-Pearson DL. Estrogen replacement therapy in the patient treated for endometrial cancer. *Obstet Gynecol* 1986;67:326-330. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3003636>.

248. Lee RB, Burke TW, Park RC. Estrogen replacement therapy following treatment for stage I endometrial carcinoma. *Gynecol Oncol* 1990;36:189-191. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2298408>.

249. Chapman JA, DiSaia PJ, Osann K, et al. Estrogen replacement in surgical stage I and II endometrial cancer survivors. *Am J Obstet Gynecol* 1996;175:1195-1200. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8942487>.

250. Barakat RR, Bundy BN, Spirtos NM, et al. Randomized double-blind trial of estrogen replacement therapy versus placebo in stage I or II endometrial cancer: a Gynecologic Oncology Group Study. *J Clin Oncol* 2006;24:587-592. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16446331>.

251. Chlebowski RT, Hendrix SL, Langer RD, et al. Influence of estrogen plus progestin on breast cancer and mammography in healthy postmenopausal women: the Women's Health Initiative Randomized Trial. *JAMA* 2003;289:3243-3253. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12824205>.

252. Anderson GL, Limacher M, Assaf AR, et al. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA* 2004;291:1701-1712. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15082697>.

253. LaCroix AZ, Chlebowski RT, Manson JE, et al. Health outcomes after stopping conjugated equine estrogens among postmenopausal women with prior hysterectomy: a randomized controlled trial. *JAMA* 2011;305:1305-1314. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21467283>.

254. Pachman DR, Jones JM, Loprinzi CL. Management of menopause-associated vasomotor symptoms: Current treatment options, challenges and future directions. *Int J Womens Health* 2010;2:123-135. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21072305>.

255. Morrow PK, Mattair DN, Hortobagyi GN. Hot flashes: a review of pathophysiology and treatment modalities. *Oncologist* 2011;16:1658-1664. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22042786>.

256. Levine JP. Treating menopausal symptoms with a tissue-selective estrogen complex. *Gend Med* 2011;8:57-68. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21536225>.

257. Pinkerton JV, Utian WH, Constantine GD, et al. Relief of vasomotor symptoms with the tissue-selective estrogen complex containing bazedoxifene/conjugated estrogens: a randomized, controlled trial. *Menopause* 2009;16:1116-1124. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19546826>.

258. Loprinzi CL, Barton DL, Qin R. Nonestrogenic management of hot flashes. *J Clin Oncol* 2011;29:3842-3846. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21911722>.

259. Sideras K, Loprinzi CL. Nonhormonal management of hot flashes for women on risk reduction therapy. *J Natl Compr Canc Netw* 2010;8:1171-1179. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20971841>.

260. Jhingran A, Burke TW, Eifel PJ. Definitive radiotherapy for patients with isolated vaginal recurrence of endometrial carcinoma after hysterectomy. *Int J Radiat Oncol Biol Phys* 2003;56:1366-1372. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12873682>.

261. Creutzberg CL, van Putten WL, Koper PC, et al. Survival after relapse in patients with endometrial cancer: results from a randomized trial. *Gynecol Oncol* 2003;89:201-209. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12713981>.

262. Lin LL, Grigsby PW, Powell MA, Mutch DG. Definitive radiotherapy in the management of isolated vaginal recurrences of endometrial cancer. *Int J Radiat Oncol Biol Phys* 2005;63:500-504. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16168841>.

263. Barakat RR, Goldman NA, Patel DA, et al. Pelvic exenteration for recurrent endometrial cancer. *Gynecol Oncol* 1999;75:99-102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10502433>.

264. Fleisch MC, Pantke P, Beckmann MW, et al. Predictors for long-term survival after interdisciplinary salvage surgery for advanced or recurrent gynecologic cancers. *J Surg Oncol* 2007;95:476-484. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17192947>.

265. Dowdy SC, Mariani A, Cliby WA, et al. Radical pelvic resection and intraoperative radiation therapy for recurrent endometrial cancer: technique and analysis of outcomes. *Gynecol Oncol* 2006;101:280-286. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16321431>.

266. Tran PT, Su Z, Hara W, et al. Long-term survivors using intraoperative radiotherapy for recurrent gynecologic malignancies. *Int J Radiat Oncol Biol Phys* 2007;69:504-511. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17560736>.

267. Whitney CW, Brunetto VL, Zaino RJ, et al. Phase II study of medroxyprogesterone acetate plus tamoxifen in advanced endometrial carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol* 2004;92:4-9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14751130>.

268. Herzog TJ. What is the clinical value of adding tamoxifen to progestins in the treatment [correction for treatment] of advanced or recurrent endometrial cancer? *Gynecol Oncol* 2004;92:1-3. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14751129>.

269. Singh M, Zaino RJ, Filiaci VJ, Leslie KK. Relationship of estrogen and progesterone receptors to clinical outcome in metastatic endometrial carcinoma: a Gynecologic Oncology Group Study. *Gynecol Oncol* 2007;106:325-333. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17532033>.

270. Kauppila A. Oestrogen and progestin receptors as prognostic indicators in endometrial cancer. A review of the literature. *Acta Oncol* 1989;28:561-566. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2675940>.

271. Thigpen JT, Brady MF, Alvarez RD, et al. Oral medroxyprogesterone acetate in the treatment of advanced or recurrent endometrial carcinoma: a dose-response study by the Gynecologic Oncology Group. *J Clin Oncol* 1999;17:1736-1744. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10561210>.

272. Dellinger TH, Monk BJ. Systemic therapy for recurrent endometrial cancer: a review of North American trials. *Expert Rev Anticancer Ther* 2009;9:905-916. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19589030>.

273. Quinn MA, Campbell JJ. Tamoxifen therapy in advanced/recurrent endometrial carcinoma. *Gynecol Oncol* 1989;32:1-3. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2909443>.

274. Thigpen T, Brady MF, Homesley HD, et al. Tamoxifen in the treatment of advanced or recurrent endometrial carcinoma: a Gynecologic Oncology Group study. *J Clin Oncol* 2001;19:364-367. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11208827>.

275. Pandya KJ, Yeap BY, Weiner LM, et al. Megestrol and tamoxifen in patients with advanced endometrial cancer: an Eastern Cooperative

Oncology Group Study (E4882). *Am J Clin Oncol* 2001;24:43-46. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11232948>.

276. McMeekin DS, Gordon A, Fowler J, et al. A phase II trial of arzoxifene, a selective estrogen response modulator, in patients with recurrent or advanced endometrial cancer. *Gynecol Oncol* 2003;90:64-69. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12821343>.

277. Quinn MA. Hormonal treatment of endometrial cancer. *Hematol Oncol Clin North Am* 1999;13:163-187. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10080075>.

278. Ray M, Fleming G. Management of advanced-stage and recurrent endometrial cancer. *Semin Oncol* 2009;36:145-154. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19332249>.

279. Humber CE, Tierney JF, Symonds RP, et al. Chemotherapy for advanced, recurrent or metastatic endometrial cancer: a systematic review of Cochrane collaboration. *Ann Oncol* 2007;18:409-420. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17150999>.

280. Cella D, Huang H, Homesley HD, et al. Patient-reported peripheral neuropathy of doxorubicin and cisplatin with and without paclitaxel in the treatment of advanced endometrial cancer: Results from GOG 184. *Gynecol Oncol* 2010;119:538-542. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20863554>.

281. Fleming GF, Brunetto VL, Cella D, et al. Phase III trial of doxorubicin plus cisplatin with or without paclitaxel plus filgrastim in advanced endometrial carcinoma: a Gynecologic Oncology Group Study. *J Clin Oncol* 2004;22:2159-2166. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15169803>.

282. Miller D, Filiaci V, Fleming G, et al. Randomized phase III noninferiority trial of first line chemotherapy for metastatic or recurrent endometrial carcinoma: A Gynecologic Oncology Group study [abstract]. *Gynecologic Oncology* 2012;125:771. Available at: [http://www.gynecologiconcology-online.net/article/S0090-8258\(12\)00228-4/abstract](http://www.gynecologiconcology-online.net/article/S0090-8258(12)00228-4/abstract).

283. Sovak MA, Dupont J, Hensley ML, et al. Paclitaxel and carboplatin in the treatment of advanced or recurrent endometrial cancer: a large retrospective study. *Int J Gynecol Cancer* 2007;17:197-203. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17291253>.

284. Pectasides D, Xiros N, Papaxoinis G, et al. Carboplatin and paclitaxel in advanced or metastatic endometrial cancer. *Gynecol Oncol* 2008;109:250-254. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18299146>.

285. Sorbe B, Andersson H, Boman K, et al. Treatment of primary advanced and recurrent endometrial carcinoma with a combination of carboplatin and paclitaxel-long-term follow-up. *Int J Gynecol Cancer* 2008;18:803-808. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17944917>.

286. Nomura H, Aoki D, Takahashi F, et al. Randomized phase II study comparing docetaxel plus cisplatin, docetaxel plus carboplatin, and paclitaxel plus carboplatin in patients with advanced or recurrent endometrial carcinoma: a Japanese Gynecologic Oncology Group study (JGOG2041). *Ann Oncol* 2011;22:636-642. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20696677>.

287. Simpkins F, Drake R, Escobar PF, et al. A phase II trial of paclitaxel, carboplatin, and bevacizumab in advanced and recurrent endometrial carcinoma (EMCA). *Gynecol Oncol* 2015;136:240-245. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25485782>.

288. Rose PG, Ali S, Moslemi-Kebria M, Simpkins F. Paclitaxel, Carboplatin, and Bevacizumab in Advanced and Recurrent Endometrial Carcinoma. *Int J Gynecol Cancer* 2017;27:452-458. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28187088>.

289. Slomovitz BM, Jiang Y, Yates MS, et al. Phase II study of everolimus and letrozole in patients with recurrent endometrial carcinoma. *J Clin Oncol* 2015;33:930-936. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25624430>.

290. Wadler S, Levy DE, Lincoln ST, et al. Topotecan is an active agent in the first-line treatment of metastatic or recurrent endometrial carcinoma: Eastern Cooperative Oncology Group Study E3E93. *J Clin Oncol* 2003;21:2110-2114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12775736>.
291. Traina TA, Sabbatini P, Aghajanian C, Dupont J. Weekly topotecan for recurrent endometrial cancer: a case series and review of the literature. *Gynecol Oncol* 2004;95:235-241. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15385138>.
292. Miller DS, Blessing JA, Lentz SS, Waggoner SE. A phase II trial of topotecan in patients with advanced, persistent, or recurrent endometrial carcinoma: a gynecologic oncology group study. *Gynecol Oncol* 2002;87:247-251. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12468321>.
293. Moxley KM, McMeekin DS. Endometrial carcinoma: a review of chemotherapy, drug resistance, and the search for new agents. *Oncologist* 2010;15:1026-1033. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20930101>.
294. Muggia FM, Blessing JA, Sorosky J, Reid GC. Phase II trial of the pegylated liposomal doxorubicin in previously treated metastatic endometrial cancer: a Gynecologic Oncology Group study. *J Clin Oncol* 2002;20:2360-2364. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11981008>.
295. Garcia AA, Blessing JA, Nolte S, Mannel RS. A phase II evaluation of weekly docetaxel in the treatment of recurrent or persistent endometrial carcinoma: a study by the Gynecologic Oncology Group. *Gynecol Oncol* 2008;111:22-26. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18675446>.
296. Ott PA, Bang YJ, Berton-Rigaud D, et al. Safety and Antitumor Activity of Pembrolizumab in Advanced Programmed Death Ligand 1-Positive Endometrial Cancer: Results From the KEYNOTE-028 Study. *J Clin Oncol* 2017;35:2535-2541. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28489510>.
297. Le DT, Durham JN, Smith KN, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 2017;357:409-413. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28596308>.
298. Le DT, Uram JN, Wang H, et al. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med* 2015;372:2509-2520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26028255>.
299. Prescribing Information: Pembrolizumab 2017. Available at: <http://bit.ly/2cTmltE>. Accessed Jul 25, 2017.
300. Alvarez EA, Brady WE, Walker JL, et al. Phase II trial of combination bevacizumab and temsirolimus in the treatment of recurrent or persistent endometrial carcinoma: a Gynecologic Oncology Group study. *Gynecol Oncol* 2013;129:22-27. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23262204>.
301. Aghajanian C, Sill MW, Darcy KM, et al. Phase II trial of bevacizumab in recurrent or persistent endometrial cancer: A Gynecologic Oncology Group Study. *J Clin Oncol* 2011;29:2259-2265. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21537039>.
302. Oza AM, Elit L, Tsao MS, et al. Phase II study of temsirolimus in women with recurrent or metastatic endometrial cancer: a trial of the NCIC Clinical Trials Group. *J Clin Oncol* 2011;29:3278-3285. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21788564>.
303. Fleming GF, Filiaci VL, Marzullo B, et al. Temsirolimus with or without megestrol acetate and tamoxifen for endometrial cancer: a gynecologic oncology group study. *Gynecol Oncol* 2014;132:585-592. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24456823>.
304. Boruta DM, 2nd, Gehrig PA, Fader AN, Olawaiye AB. Management of women with uterine papillary serous cancer: a Society of Gynecologic Oncology (SGO) review. *Gynecol Oncol* 2009;115:142-153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19592079>.

305. Olawaiye AB, Boruta DM, 2nd. Management of women with clear cell endometrial cancer: a Society of Gynecologic Oncology (SGO) review. *Gynecol Oncol* 2009;113:277-283. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19251307>.
306. Mendivil A, Schuler KM, Gehrig PA. Non-endometrioid adenocarcinoma of the uterine corpus: a review of selected histological subtypes. *Cancer Control* 2009;16:46-52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19078929>.
307. Varughese J, Hui P, Lu L, et al. Clear cell cancer of the uterine corpus: the association of clinicopathologic parameters and treatment on disease progression. *J Oncol* 2011;2011:628084. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22187554>.
308. Kernochan LE, Garcia RL. Carcinosarcomas (malignant mixed Mullerian tumor) of the uterus: advances in elucidation of biologic and clinical characteristics. *J Natl Compr Canc Netw* 2009;7:550-556; quiz 557. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19460280>.
309. Cirisano FD, Jr., Robboy SJ, Dodge RK, et al. The outcome of stage I-II clinically and surgically staged papillary serous and clear cell endometrial cancers when compared with endometrioid carcinoma. *Gynecol Oncol* 2000;77:55-65. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10739691>.
310. Hendrickson MR, Longacre TA, Kempson RL. Uterine papillary serous carcinoma revisited. *Gynecol Oncol* 1994;54:261-263. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8088601>.
311. McCluggage WG. Uterine carcinosarcomas (malignant mixed Mullerian tumors) are metaplastic carcinomas. *Int J Gynecol Cancer* 2002;12:687-690. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12445244>.
312. Cantrell LA, Havrilesky L, Moore DT, et al. A multi-institutional cohort study of adjuvant therapy in stage I-II uterine carcinosarcoma. *Gynecol Oncol* 2012;127:22-26. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22727985>.
313. Kanthan R, Senger JL. Uterine carcinosarcomas (malignant mixed mullerian tumours): a review with special emphasis on the controversies in management. *Obstet Gynecol Int* 2011;2011:470795. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22007228>.
314. D'Angelo E, Prat J. Pathology of mixed Mullerian tumours. *Best Pract Res Clin Obstet Gynaecol* 2011;25:705-718. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21742560>.
315. de Jong RA, Nijman HW, Wijbrandi TF, et al. Molecular markers and clinical behavior of uterine carcinosarcomas: focus on the epithelial tumor component. *Mod Pathol* 2011;24:1368-1379. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21572397>.
316. Vorgias G, Fotiou S. The role of lymphadenectomy in uterine carcinosarcomas (malignant mixed mullerian tumours): a critical literature review. *Arch Gynecol Obstet* 2010;282:659-664. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20721670>.
317. Vogel TJ, Knickerbocker A, Shah CA, et al. An analysis of current treatment practice in uterine papillary serous and clear cell carcinoma at two high volume cancer centers. *J Gynecol Oncol* 2015;26:25-31. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25376917>.
318. Monterossi G, Ghezzi F, Vizza E, et al. Minimally Invasive Approach in Type II Endometrial Cancer: Is It Wise and Safe? *J Minim Invasive Gynecol* 2017;24:438-445. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28065812>.
319. Fader AN, Seamon LG, Escobar PF, et al. Minimally invasive surgery versus laparotomy in women with high grade endometrial cancer: a multi-site study performed at high volume cancer centers. *Gynecol Oncol* 2012;126:180-185. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22555102>.
320. Koskas M, Jozwiak M, Fournier M, et al. Long-term oncological safety of minimally invasive surgery in high-risk endometrial cancer. *Eur J Cancer* 2016;65:185-191. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27505629>.

321. Salehi S, Avall-Lundqvist E, Legerstam B, et al. Robot-assisted laparoscopy versus laparotomy for infrarenal paraaortic lymphadenectomy in women with high-risk endometrial cancer: A randomised controlled trial. *Eur J Cancer* 2017;79:81-89. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28463759>.

322. Fader AN, Drake RD, O'Malley DM, et al. Platinum/taxane-based chemotherapy with or without radiation therapy favorably impacts survival outcomes in stage I uterine papillary serous carcinoma. *Cancer* 2009;115:2119-2127. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19306417>.

323. Fader AN, Nagel C, Axtell AE, et al. Stage II uterine papillary serous carcinoma: Carboplatin/paclitaxel chemotherapy improves recurrence and survival outcomes. *Gynecol Oncol* 2009;112:558-562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19118888>.

324. Vandenput I, Trovik J, Vergote I, et al. The role of adjuvant chemotherapy in surgical stages I-II serous and clear cell carcinomas and carcinosarcoma of the endometrium: A collaborative study. *Int J Gynecol Cancer* 2011;21:332-336. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21270614>.

325. Kelly MG, O'Malley D M, Hui P, et al. Improved survival in surgical stage I patients with uterine papillary serous carcinoma (UPSC) treated with adjuvant platinum-based chemotherapy. *Gynecol Oncol* 2005;98:353-359. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16005947>.

326. Thomas MB, Mariani A, Cliby WA, et al. Role of systematic lymphadenectomy and adjuvant therapy in stage I uterine papillary serous carcinoma. *Gynecol Oncol* 2007;107:186-189. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17688926>.

327. Goldberg H, Miller RC, Abdah-Bortnyak R, et al. Outcome after combined modality treatment for uterine papillary serous carcinoma: a study by the Rare Cancer Network (RCN). *Gynecol Oncol* 2008;108:298-305. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18096209>.

328. Hamilton CA, Cheung MK, Osann K, et al. The effect of adjuvant chemotherapy versus whole abdominopelvic radiation on the survival of patients with advanced stage uterine papillary serous carcinoma. *Gynecol Oncol* 2006;103:679-683. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16793126>.

329. Grice J, Ek M, Greer B, et al. Uterine papillary serous carcinoma: evaluation of long-term survival in surgically staged patients. *Gynecol Oncol* 1998;69:69-73. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9571001>.

330. Havrilesky LJ, Secord AA, Bae-Jump V, et al. Outcomes in surgical stage I uterine papillary serous carcinoma. *Gynecol Oncol* 2007;105:677-682. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17355889>.

331. Velker V, D'Souza D, Prefontaine M, et al. Role of Adjuvant Therapy for Stage IA Serous and Clear Cell Uterine Cancer: Is Observation a Valid Strategy? *Int J Gynecol Cancer* 2016;26:491-496. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26825823>.

332. Sutton G, Axelrod JH, Bundy BN, et al. Adjuvant whole abdominal irradiation in clinical stages I and II papillary serous or clear cell carcinoma of the endometrium: a phase II study of the Gynecologic Oncology Group. *Gynecol Oncol* 2006;100:349-354. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16213007>.

333. Fader AN, Boruta D, Olawaiye AB, Gehrig PA. Uterine papillary serous carcinoma: epidemiology, pathogenesis and management. *Curr Opin Obstet Gynecol* 2010;22:21-29. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19952744>.

334. Homesley HD, Filiaci V, Markman M, et al. Phase III trial of ifosfamide with or without paclitaxel in advanced uterine carcinosarcoma: a Gynecologic Oncology Group Study. *J Clin Oncol* 2007;25:526-531. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17290061>.

335. Einstein MH, Frimer M, Kuo DY, et al. Phase II trial of adjuvant pelvic radiation "sandwiched" between combination paclitaxel and carboplatin in women with uterine papillary serous carcinoma. *Gynecol Oncol*

2012;124:21-25. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22035806>.

336. Galaal K, van der Heijden E, Godfrey K, et al. Adjuvant radiotherapy and/or chemotherapy after surgery for uterine carcinosarcoma. *Cochrane Database Syst Rev* 2013;2:CD006812. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23450572>.

337. Wang W, Do V, Hogg R, et al. Uterine papillary serous carcinoma: patterns of failure and survival. *Aust N Z J Obstet Gynaecol* 2009;49:419-425. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19694700>.

338. Mehta N, Yamada SD, Rotmensch J, Mundt AJ. Outcome and pattern of failure in pathologic stage I-II papillary serous carcinoma of the endometrium: implications for adjuvant radiation therapy. *Int J Radiat Oncol Biol Phys* 2003;57:1004-1009. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/14575831>.

339. Murphy KT, Rotmensch J, Yamada SD, Mundt AJ. Outcome and patterns of failure in pathologic stages I-IV clear-cell carcinoma of the endometrium: implications for adjuvant radiation therapy. *Int J Radiat Oncol Biol Phys* 2003;55:1272-1276. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12654437>.

340. Sood BM, Jones J, Gupta S, et al. Patterns of failure after the multimodality treatment of uterine papillary serous carcinoma. *Int J Radiat Oncol Biol Phys* 2003;57:208-216. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12909235>.

341. Wolfson AH, Brady MF, Rocereto T, et al. A gynecologic oncology group randomized phase III trial of whole abdominal irradiation (WAI) vs. cisplatin-ifosfamide and mesna (CIM) as post-surgical therapy in stage I-IV carcinosarcoma (CS) of the uterus. *Gynecol Oncol* 2007;107:177-185. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17822748>.

342. Jhingran A, Ramondetta LM, Bodurka DC, et al. A prospective phase II study of chemoradiation followed by adjuvant chemotherapy for FIGO stage I-IIIa (1988) uterine papillary serous carcinoma of the endometrium.

*Gynecol Oncol* 2013;129:304-309. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23385150>.

343. Mahdi H, Rose PG, Elshaikh MA, et al. Adjuvant vaginal brachytherapy decreases the risk of vaginal recurrence in patients with stage I non-invasive uterine papillary serous carcinoma. A multi-institutional study. *Gynecol Oncol* 2015;136:529-533. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25575483>.

344. Mahdi H, Elshaikh MA, DeBenardo R, et al. Impact of adjuvant chemotherapy and pelvic radiation on pattern of recurrence and outcome in stage I non-invasive uterine papillary serous carcinoma. A multi-institution study. *Gynecol Oncol* 2015;137:239-244. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25641568>.

345. Fader AN, Roque DM, Siegel E, et al. Randomized Phase II Trial of Carboplatin-Paclitaxel Versus Carboplatin-Paclitaxel-Trastuzumab in Uterine Serous Carcinomas That Overexpress Human Epidermal Growth Factor Receptor 2/neu. *J Clin Oncol* 2018;JCO2017765966. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29584549>.

346. Sutton G, Brunetto VL, Kilgore L, et al. A phase III trial of ifosfamide with or without cisplatin in carcinosarcoma of the uterus: A Gynecologic Oncology Group Study. *Gynecol Oncol* 2000;79:147-153. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11063636>.

347. Hensley ML. Role of chemotherapy and biomolecular therapy in the treatment of uterine sarcomas. *Best Pract Res Clin Obstet Gynaecol* 2011;25:773-782. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21752717>.

348. Sutton G, Kauderer J, Carson LF, et al. Adjuvant ifosfamide and cisplatin in patients with completely resected stage I or II carcinosarcomas (mixed mesodermal tumors) of the uterus: a Gynecologic Oncology Group study. *Gynecol Oncol* 2005;96:630-634. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15721404>.

349. Powell MA, Filiaci VL, Rose PG, et al. Phase II evaluation of paclitaxel and carboplatin in the treatment of carcinosarcoma of the

uterus: a Gynecologic Oncology Group study. *J Clin Oncol* 2010;28:2727-2731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20421537>.

350. Reed NS, Mangioni C, Malmström H, et al. Phase III randomised study to evaluate the role of adjuvant pelvic radiotherapy in the treatment of uterine sarcomas stages I and II: an European Organisation for Research and Treatment of Cancer Gynaecological Cancer Group Study (protocol 55874). *Eur J Cancer* 2008;44:808-818. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18378136>.

351. Callister M, Ramondetta LM, Jhingran A, et al. Malignant mixed Müllerian tumors of the uterus: analysis of patterns of failure, prognostic factors, and treatment outcome. *Int J Radiat Oncol Biol Phys* 2004;58:786-796. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14967435>.

352. Chi DS, Mychalczak B, Saigo PE, et al. The role of whole-pelvic irradiation in the treatment of early-stage uterine carcinosarcoma. *Gynecol Oncol* 1997;65:493-498. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9190981>.

353. Knocke TH, Weitmann HD, Kucera H, et al. Results of primary and adjuvant radiotherapy in the treatment of mixed Müllerian tumors of the corpus uteri. *Gynecol Oncol* 1999;73:389-395. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10366465>.

354. Larson B, Silfverswärd C, Nilsson B, Pettersson F. Mixed müllerian tumours of the uterus--prognostic factors: a clinical and histopathologic study of 147 cases. *Radiother Oncol* 1990;17:123-132. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2157241>.

355. Gerszten K, Faul C, Kounelis S, et al. The impact of adjuvant radiotherapy on carcinosarcoma of the uterus. *Gynecol Oncol* 1998;68:8-13. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9454652>.

356. Dusenbery KE, Potish RA, Argenta PA, Judson PL. On the apparent failure of adjuvant pelvic radiotherapy to improve survival for women with uterine sarcomas confined to the uterus. *Am J Clin Oncol* 2005;28:295-300. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15923804>.

357. Nemani D, Mitra N, Guo M, Lin L. Assessing the effects of lymphadenectomy and radiation therapy in patients with uterine carcinosarcoma: a SEER analysis. *Gynecol Oncol* 2008;111:82-88. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18674808>.

358. Cancer Facts and Figures 2017. Atlanta, GA: American Cancer Society. Available at: <https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/cancer-facts-figures-2017.html>.

359. Trope CG, Abeler VM, Kristensen GB. Diagnosis and treatment of sarcoma of the uterus. A review. *Acta Oncol* 2012;51:694-705. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22793037>.

360. Amant F, Coosemans A, Debiec-Rychter M, et al. Clinical management of uterine sarcomas. *Lancet Oncol* 2009;10:1188-1198. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19959075>.

361. Novetsky AP, Powell MA. Management of sarcomas of the uterus. *Curr Opin Oncol* 2013;25:546-552. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23942299>.

362. Lax SF. Molecular genetic changes in epithelial, stromal and mixed neoplasms of the endometrium. *Pathology* 2007;39:46-54. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17365822>.

363. Huang HY, Ladanyi M, Soslow RA. Molecular detection of JAZF1-JJAZ1 gene fusion in endometrial stromal neoplasms with classic and variant histology: evidence for genetic heterogeneity. *Am J Surg Pathol* 2004;28:224-232. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15043312>.

364. Leath CA, 3rd, Huh WK, Hyde J, Jr., et al. A multi-institutional review of outcomes of endometrial stromal sarcoma. *Gynecol Oncol* 2007;105:630-634. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17320937>.

365. Koontz JI, Soreng AL, Nucci M, et al. Frequent fusion of the JAZF1 and JJAZ1 genes in endometrial stromal tumors. *Proc Natl Acad Sci U S A*

2001;98:6348-6353. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11371647>.

366. Lee CH, Marino-Enriquez A, Ou W, et al. The clinicopathologic features of YWHAЕ-FAM22 endometrial stromal sarcomas: a histologically high-grade and clinically aggressive tumor. *Am J Surg Pathol* 2012;36:641-653. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22456610>.

367. Sciallis AP, Bedroske PP, Schoolmeester JK, et al. High-grade endometrial stromal sarcomas: a clinicopathologic study of a group of tumors with heterogenous morphologic and genetic features. *Am J Surg Pathol* 2014;38:1161-1172. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25133706>.

368. Kurman RJ, Carcangiu ML, Herrington CS, Young RH. WHO Classification of Tumours of Female Reproductive Organs. Fourth Edition. Vol. 6: World Health Organization; 2014.

369. Zivanovic O, Leitao MM, Iasonos A, et al. Stage-specific outcomes of patients with uterine leiomyosarcoma: a comparison of the international Federation of gynecology and obstetrics and american joint committee on cancer staging systems. *J Clin Oncol* 2009;27:2066-2072. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19255317>.

370. Prat J. FIGO staging for uterine sarcomas. *Int J Gynaecol Obstet* 2009;104:177-178. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19135669>.

371. Rauh-Hain JA, del Carmen MG. Endometrial stromal sarcoma: a systematic review. *Obstet Gynecol* 2013;122:676-683. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23921879>.

372. Group EESNW. Soft tissue and visceral sarcomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2012;23 Suppl 7:vii92-99. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22997462>.

373. Barney B, Tward JD, Skidmore T, Gaffney DK. Does radiotherapy or lymphadenectomy improve survival in endometrial stromal sarcoma? *Int J Gynecol Cancer* 2009;19:1232-1238. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19823060>.

374. Shah JP, Bryant CS, Kumar S, et al. Lymphadenectomy and ovarian preservation in low-grade endometrial stromal sarcoma. *Obstet Gynecol* 2008;112:1102-1108. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18978112>.

375. Signorelli M, Fruscio R, Dell'Anna T, et al. Lymphadenectomy in uterine low-grade endometrial stromal sarcoma: an analysis of 19 cases and a literature review. *Int J Gynecol Cancer* 2010;20:1363-1366. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21051978>.

376. Reichardt P. The treatment of uterine sarcomas. *Ann Oncol* 2012;23 Suppl 10:x151-157. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22987952>.

377. Thanopoulou E, Judson I. Hormonal therapy in gynecological sarcomas. *Expert Rev Anticancer Ther* 2012;12:885-894. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22845404>.

378. Pink D, Lindner T, Mrozek A, et al. Harm or benefit of hormonal treatment in metastatic low-grade endometrial stromal sarcoma: single center experience with 10 cases and review of the literature. *Gynecol Oncol* 2006;101:464-469. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16368128>.

379. Reich O, Regauer S. Estrogen replacement therapy and tamoxifen are contraindicated in patients with endometrial stromal sarcoma. *Gynecol Oncol* 2006;102:413-414; author reply 414. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16712906>.

380. Mansi JL, Ramachandra S, Wiltshaw E, Fisher C. Endometrial stromal sarcomas. *Gynecol Oncol* 1990;36:113-118. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/2295442>.

381. Berchuck A, Rubin SC, Hoskins WJ, et al. Treatment of endometrial stromal tumors. *Gynecol Oncol* 1990;36:60-65. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2295453>.

382. Weitmann HD, Knocke TH, Kucera H, Pötter R. Radiation therapy in the treatment of endometrial stromal sarcoma. *Int J Radiat Oncol Biol Phys* 2001;49:739-748. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11172957>.

383. Brenner DJ, Hall EJ. Computed tomography--an increasing source of radiation exposure. *N Engl J Med* 2007;357:2277-2284. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18046031>.

384. Sampath S, Schultheiss TE, Ryu JK, Wong JY. The role of adjuvant radiation in uterine sarcomas. *Int J Radiat Oncol Biol Phys* 2010;76:728-734. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19700247>.

385. Mahdavi A, Monk BJ, Ragazzo J, et al. Pelvic radiation improves local control after hysterectomy for uterine leiomyosarcoma: a 20-year experience. *Int J Gynecol Cancer* 2009;19:1080-1084. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19820372>.

386. Giuntoli RL, 2nd, Metzinger DS, DiMarco CS, et al. Retrospective review of 208 patients with leiomyosarcoma of the uterus: prognostic indicators, surgical management, and adjuvant therapy. *Gynecol Oncol* 2003;89:460-469. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12798712>.

387. Dusenbery KE, Potish RA, Judson P. Limitations of adjuvant radiotherapy for uterine sarcomas spread beyond the uterus. *Gynecol Oncol* 2004;94:191-196. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15262141>.

388. Ricci S, Giuntoli RL, 2nd, Eisenhauer E, et al. Does adjuvant chemotherapy improve survival for women with early-stage uterine leiomyosarcoma? *Gynecol Oncol* 2013;131:629-633. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24016408>.

389. Hensley ML, Wathen JK, Maki RG, et al. Adjuvant therapy for high-grade, uterus-limited leiomyosarcoma: results of a phase 2 trial (SARC 005). *Cancer* 2013;119:1555-1561. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23335221>.

390. Maki RG, Wathen JK, Patel SR, et al. Randomized phase II study of gemcitabine and docetaxel compared with gemcitabine alone in patients with metastatic soft tissue sarcomas: results of sarcoma alliance for research through collaboration study 002 [corrected]. *J Clin Oncol* 2007;25:2755-2763. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17602081>.

391. Davis EJ, Chugh R, Zhao L, et al. A randomised, open-label, phase II study of neo/adjuvant doxorubicin and ifosfamide versus gemcitabine and docetaxel in patients with localised, high-risk, soft tissue sarcoma. *Eur J Cancer* 2015;51:1794-1802. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26066736>.

392. Hensley ML, Ishill N, Soslow R, et al. Adjuvant gemcitabine plus docetaxel for completely resected stages I-IV high grade uterine leiomyosarcoma: Results of a prospective study. *Gynecol Oncol* 2009;112:563-567. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19135708>.

393. Hensley ML, Miller A, O'Malley DM, et al. Randomized phase III trial of gemcitabine plus docetaxel plus bevacizumab or placebo as first-line treatment for metastatic uterine leiomyosarcoma: an NRG Oncology/Gynecologic Oncology Group study. *J Clin Oncol* 2015;33:1180-1185. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25713428>.

394. Hensley ML, Maki R, Venkatraman E, et al. Gemcitabine and docetaxel in patients with unresectable leiomyosarcoma: results of a phase II trial. *J Clin Oncol* 2002;20:2824-2831. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12065559>.

395. Karavasilis V, Seddon BM, Ashley S, et al. Significant clinical benefit of first-line palliative chemotherapy in advanced soft-tissue sarcoma: retrospective analysis and identification of prognostic factors in 488

patients. *Cancer* 2008;112:1585-1591. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/18278813>.

396. Garcia-Del-Muro X, Lopez-Pousa A, Maurel J, et al. Randomized phase II study comparing gemcitabine plus dacarbazine versus dacarbazine alone in patients with previously treated soft tissue sarcoma: a Spanish Group for Research on Sarcomas study. *J Clin Oncol* 2011;29:2528-2533. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/21606430>.

397. van der Graaf WT, Blay JY, Chawla SP, et al. Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet* 2012;379:1879-1886. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/22595799>.

398. Mancari R, Signorelli M, Gadducci A, et al. Adjuvant chemotherapy in stage I-II uterine leiomyosarcoma: a multicentric retrospective study of 140 patients. *Gynecol Oncol* 2014;133:531-536. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/24631454>.

399. Gottlieb JA, Benjamin RS, Baker LH, et al. Role of DTIC (NSC-45388) in the chemotherapy of sarcomas. *Cancer Treat Rep* 1976;60:199-203. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/769974>.

400. Sleijfer S, Ray-Coquard I, Papai Z, et al. Pazopanib, a multikinase angiogenesis inhibitor, in patients with relapsed or refractory advanced soft tissue sarcoma: a phase II study from the European organisation for research and treatment of cancer-soft tissue and bone sarcoma group (EORTC study 62043). *J Clin Oncol* 2009;27:3126-3132. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/19451427>.

401. Somaiah N, von Mehren M. New drugs and combinations for the treatment of soft-tissue sarcoma: a review. *Cancer Manag Res* 2012;4:397-411. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/23226072>.

402. Rajendra R, Jones RL, Pollack SM. Targeted treatment for advanced soft tissue sarcoma: profile of pazopanib. *Onco Targets Ther* 2013;6:217-222. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23524973>.

403. Pautier P, Floquet A, Penel N, et al. Randomized multicenter and stratified phase II study of gemcitabine alone versus gemcitabine and docetaxel in patients with metastatic or relapsed leiomyosarcomas: a Federation Nationale des Centres de Lutte Contre le Cancer (FNCLCC) French Sarcoma Group Study (TAXOGEM study). *Oncologist* 2012;17:1213-1220. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/22907974>.

404. van Hoesel QG, Verweij J, Catimel G, et al. Phase II study with docetaxel (Taxotere) in advanced soft tissue sarcomas of the adult. EORTC Soft Tissue and Bone Sarcoma Group. *Ann Oncol* 1994;5:539-542. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7918126>.

405. Edmonson JH, Ebbert LP, Nascimento AG, et al. Phase II study of docetaxel in advanced soft tissue sarcomas. *Am J Clin Oncol* 1996;19:574-576. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/8931674>.

406. Ridolfi C, Pasini G, Drudi F, et al. Long lasting clinical response to chemotherapy for advanced uterine leiomyosarcoma: a case report. *J Med Case Rep* 2013;7:29. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/23347560>.

407. Ferriss JS, Atkins KA, Lachance JA, et al. Temozolomide in advanced and recurrent uterine leiomyosarcoma and correlation with o6-methylguanine DNA methyltransferase expression: a case series. *Int J Gynecol Cancer* 2010;20:120-125. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/20130512>.

408. Anderson S, Aghajanian C. Temozolomide in uterine leiomyosarcomas. *Gynecol Oncol* 2005;98:99-103. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/15916799>.

409. Talbot SM, Keohan ML, Hesdorffer M, et al. A phase II trial of temozolomide in patients with unresectable or metastatic soft tissue sarcoma. *Cancer* 2003;98:1942-1946. Available at:  
<http://www.ncbi.nlm.nih.gov/pubmed/14584078>.

410. Oosten AW, Seynaeve C, Schmitz PI, et al. Outcomes of first-line chemotherapy in patients with advanced or metastatic leiomyosarcoma of uterine and non-uterine origin. *Sarcoma* 2009;2009:348910. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20066161>.
411. Bernstein-Molho R, Grisaro D, Soyfer V, et al. Metastatic uterine leiomyosarcomas: a single-institution experience. *Int J Gynecol Cancer* 2010;20:255-260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20134269>.
412. Look KY, Sandler A, Blessing JA, et al. Phase II trial of gemcitabine as second-line chemotherapy of uterine leiomyosarcoma: a Gynecologic Oncology Group (GOG) Study. *Gynecol Oncol* 2004;92:644-647. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14766260>.
413. Judson I, Radford JA, Harris M, et al. Randomised phase II trial of pegylated liposomal doxorubicin (DOXIL/CAELYX) versus doxorubicin in the treatment of advanced or metastatic soft tissue sarcoma: a study by the EORTC Soft Tissue and Bone Sarcoma Group. *Eur J Cancer* 2001;37:870-877. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11313175>.
414. Sutton G, Blessing J, Hanjani P, Kramer P. Phase II evaluation of liposomal doxorubicin (Doxil) in recurrent or advanced leiomyosarcoma of the uterus: a Gynecologic Oncology Group study. *Gynecol Oncol* 2005;96:749-752. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15721421>.
415. Gallup DG, Blessing JA, Andersen W, Morgan MA. Evaluation of paclitaxel in previously treated leiomyosarcoma of the uterus: a gynecologic oncology group study. *Gynecol Oncol* 2003;89:48-51. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12694653>.
416. George S, Feng Y, Manola J, et al. Phase 2 trial of aromatase inhibition with letrozole in patients with uterine leiomyosarcomas expressing estrogen and/or progesterone receptors. *Cancer* 2014;120:738-743. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24222211>.
417. Gajdos C, Elias A. Trabectedin: safety and efficacy in the treatment of advanced sarcoma. *Clin Med Insights Oncol* 2011;5:35-43. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21499557>.
418. Demetri GD, Chawla SP, von Mehren M, et al. Efficacy and safety of trabectedin in patients with advanced or metastatic liposarcoma or leiomyosarcoma after failure of prior anthracyclines and ifosfamide: results of a randomized phase II study of two different schedules. *J Clin Oncol* 2009;27:4188-4196. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19652065>.
419. Fayette J, Boyle H, Chabaud S, et al. Efficacy of trabectedin for advanced sarcomas in clinical trials versus compassionate use programs: analysis of 92 patients treated in a single institution. *Anticancer Drugs* 2010;21:113-119. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19887935>.
420. Pautier P, Floquet A, Chevreau C, et al. Trabectedin in combination with doxorubicin for first-line treatment of advanced uterine or soft-tissue leiomyosarcoma (LMS-02): a non-randomised, multicentre, phase 2 trial. *Lancet Oncol* 2015;16:457-464. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25795402>.
421. Demetri GD, von Mehren M, Jones RL, et al. Efficacy and safety of trabectedin or dacarbazine for metastatic liposarcoma or leiomyosarcoma after failure of conventional chemotherapy: results of a phase III randomized multicenter clinical trial. *J Clin Oncol* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26371143>.
422. Hensley ML, Patel SR, von Mehren M, et al. Efficacy and safety of trabectedin or dacarbazine in patients with advanced uterine leiomyosarcoma after failure of anthracycline-based chemotherapy: Subgroup analysis of a phase 3, randomized clinical trial. *Gynecol Oncol* 2017;146:531-537. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28651804>.
423. Schoffski P, Maki RG, Italiano A, et al. Randomized, open-label, multicenter, phase III study of eribulin versus dacarbazine in patients (pts) with leiomyosarcoma (LMS) and adipocytic sarcoma (ADI) [abstract].

ASCO Meeting Abstracts 2015;33:LBA10502. Available at:  
[http://meeting.ascopubs.org/cgi/content/abstract/33/18\\_suppl/LBA10502](http://meeting.ascopubs.org/cgi/content/abstract/33/18_suppl/LBA10502).

424. Yoon A, Park JY, Park JY, et al. Prognostic factors and outcomes in endometrial stromal sarcoma with the 2009 FIGO staging system: a multicenter review of 114 cases. *Gynecol Oncol* 2014;132:70-75. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24184602>.

425. Leitao MM, Brennan MF, Hensley M, et al. Surgical resection of pulmonary and extrapulmonary recurrences of uterine leiomyosarcoma. *Gynecol Oncol* 2002;87:287-294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12468327>.

426. Sharma S, Takyar S, Manson SC, et al. Efficacy and safety of pharmacological interventions in second- or later-line treatment of patients with advanced soft tissue sarcoma: a systematic review. *BMC Cancer* 2013;13:385. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23937858>.

427. Dhakal S, Corbin KS, Milano MT, et al. Stereotactic body radiotherapy for pulmonary metastases from soft-tissue sarcomas: excellent local lesion control and improved patient survival. *Int J Radiat Oncol Biol Phys* 2012;82:940-945. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21277105>.

428. Mehta N, Selch M, Wang PC, et al. Safety and efficacy of stereotactic body radiation therapy in the treatment of pulmonary metastases from high grade sarcoma. *Sarcoma* 2013;2013:360214. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24198717>.

429. Castells MC, Tennant NM, Sloane DE, et al. Hypersensitivity reactions to chemotherapy: outcomes and safety of rapid desensitization in 413 cases. *J Allergy Clin Immunol* 2008;122:574-580. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18502492>.

430. Dizon DS, Sabbatini PJ, Aghajanian C, et al. Analysis of patients with epithelial ovarian cancer or fallopian tube carcinoma retreated with cisplatin after the development of a carboplatin allergy. *Gynecol Oncol* 2002;84:378-382. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11855873>.

431. Sampson HA, Munoz-Furlong A, Campbell RL, et al. Second symposium on the definition and management of anaphylaxis: summary report--second National Institute of Allergy and Infectious Disease/Food Allergy and Anaphylaxis Network symposium. *Ann Emerg Med* 2006;47:373-380. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16546624>.

432. Manivannan V, Decker WW, Stead LG, et al. Visual representation of National Institute of Allergy and Infectious Disease and Food Allergy and Anaphylaxis Network criteria for anaphylaxis. *Int J Emerg Med* 2009;2:3-5. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19390910>.

433. Lenz HJ. Management and preparedness for infusion and hypersensitivity reactions. *Oncologist* 2007;12:601-609. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17522249>.

434. Markman M, Zanotti K, Peterson G, et al. Expanded experience with an intradermal skin test to predict for the presence or absence of carboplatin hypersensitivity. *J Clin Oncol* 2003;21:4611-4614. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14673050>.

435. Lee CW, Matulonis UA, Castells MC. Rapid inpatient/outpatient desensitization for chemotherapy hypersensitivity: standard protocol effective in 57 patients for 255 courses. *Gynecol Oncol* 2005;99:393-399. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16054201>.

436. Lee CW, Matulonis UA, Castells MC. Carboplatin hypersensitivity: a 6-h 12-step protocol effective in 35 desensitizations in patients with gynecological malignancies and mast cell/IgE-mediated reactions. *Gynecol Oncol* 2004;95:370-376. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15491759>.

437. Markman M, Hsieh F, Zanotti K, et al. Initial experience with a novel desensitization strategy for carboplatin-associated hypersensitivity reactions: carboplatin-hypersensitivity reactions. *J Cancer Res Clin Oncol* 2004;130:25-28. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14564516>.